

remota, inaequidistantia nec aggregata, elongato-lanceolata, 25-32 cent. longa, 3-4.5 cent. lata, utrinque aequaliter attenuata, acuminata, apice setoso-penicillata, supra opaca, subtus subconcolora, costulis 5 percursa, quarum medialis supra valdiuscula nuda, lateralibus minoribus et spinuloso-ciliatis, subtus nervis omnibus inconspicuis et nudis; spadicis spathae primariae elongato-tubulosae aculeolatae; spathae secundariae tubuloso-infundibuliformes inermes; spicae erecto-patulae, curvulae 4.5-6 cent. longae, floribus 10-12 distice utrinque praeditae: fructus parvi globosi 8-9 mill. diam. apiculati, perianthio pedicelliformi suffulti, squamis laxe imbricatis, semine subgloboso, superficie inaequali, albumine aequabili.

Philippines; Malanipa, (Moseley in Herb. Kew).

126. *Calamus Vidalianus* Becc. sp. n.—*Frondium rachis* in parte basilari subtus convexa et nuda, longitudinaliter ad latera sulcata, aculeis vix reduncis dense armata, in facie superiore convexiuscula dense spinosa; segmenta aequidistantia, angustissime lanceolata, basi attenuata, apici acuminata, 30-32 cent. longa, 15-20 mill. lata, glabra, utrinque subconcolora, supra costulis 3 spinulosis percursa, subtus nuda, marginibus ciliato-spinulosis; spadicis ♀ spathae primariae tubuloso-cylindraceae, in parte superiore aliquantum ampliatae, sparse aculeatae; spathae secundariae elongato-infundibuliformes inermes; spicae ad faucem spatharum callo insertae, patentia, arcuata, 5-7 cent. longa, floribus subdisticis utrinque paucis.—*C. horrens* (not of Bl.) Vidal.

Philippines (Vidal No. 933 in Herb. Kew).

127. *CALAMUS PISICARPUS* Bl.—*C. oblongus* β; Bl.—*C. verus* (not of Lour.) Mart. . . . . Moluccas

128. *CALAMUS ARUENSIS* Becc. . . . . Aru Islands.

129. *CALAMUS HOLLRUNGII* Becc. . . . . New Guinea.

130. *CALAMUS UNIFARIUS* H. Wendl. . . . . Borneo.  
*C. UNIFARIUS* v. *PENTONG* Becc. . . . . Nicobar Islands.

131. *Calamus subinermis* H. Wendl. (nomen nudum in Herb. Kew.)—Scandens, robustus, vaginis 4 cent. diam. crases lignosis, apici saccatis, inermibus, indumento tenui cinereo-furfuraceo detergibili tectis, in ore nudis et oblique truncatis; frondes magnae cirro validissimo crebre aculeis reduncis semiverticillatis armato terminatae, petiolo brevi subtus inermi; segmentis magnis 50-60 cent. longis, 5-5.5 cent. latis, non numerosis, subaequidistantibus, elongato-lanceolatis, basi longe attenuatis, apici acuminatis, utrinque sub-

concoloribus, supra costulis 5 spinuloso-setulosis percursis, subtus nudis, marginibus ciliolato-spinulosis.

Borneo (Low in Herb. Kew)

- 132. *CALAMUS ALBUS* Pers. . . . Moluccas.
- 133. *CALAMUS GRAMINOSUS* Bl. . . . Amboina.
- 134. *CALAMUS NEGLECTUS* Becc. . . . Malayan Peninsula.
- 135. *CALAMUS ASPERRIMUS* Bl. . . . Java.
- 136. *CALAMUS EQUESTRIS* Willd. . . . Moluccas.
- 137. *CALAMUS CAWA* Bl. . . . Moluccas.

138. *Calamus mucronatus* Becc. sp. n.—Scandens, gracillimus, vaginis 4-6 mill. diam. non flagelliferis inermibus vel spinis brevissimis tuberculiformibus armatis; frondes breves cirro aculeato gracillimo terminatae, petiolo subnullo; segmenta parva perpaucia remote inequidistantia alterna vel subopposita, oblanceolata, abrupte acuminato-mucronata, nerves 5 tenuibus percursa, utrinque subconcoloria et in nervis nec non in margine omnino nuda; majora 6-12 cent. longa, 12-16 mill. lata, basilaria basi callosa et reflexa; spadices breves folio breviores non flagelliferi, spathis primariis tubulosis laxe vaginantibus et sursum versus ampliatis, inermibus, spicis parvis scorpioideis.

Borneo; Sarawak (Beccari P. B. Nos. 829, 1904).

- 139. *CALAMUS MELANOLOMA* Mart.—*C. anceps* Bl.—*C. calolepis* Miq. . . . Java.
- 140. *CALAMUS VIRIDISPINUS* Becc. . . . Malayan Peninsula.
- 141. *CALAMUS EARIOACANTHUS* Becc. . . . Borneo.
- 142. *CALAMUS SIPHONOSPATHUS* Mart.—*C. discolor* Mart. —Philippines.

143. *Calamus microcarpus* Becc. sp. n.—Scandens; frondes cirriferae, rachi subtus aculeis reduncis armata supra bifaciali et in medio argute spinulosa; segmenta numerosa inaequidistantia, 3-7 approximata, linear-lanceolata, acuminatissima, 20-23 cent. longa, 12-15 mill. lata, utrinque subconcolora, supra costa media acuta et in utroque latere nervis 1-2, setis paucis rigidis obsitis, percursa, subtus et tantum costa media prope apicem setulosa, marginibus crebre et patule ciliato-spinulosis; spadices partialiter ultra decompositi, spathis tubulosis laxe vaginantibus, basi attenuatis, apici ampliatis, vix spinulosis; inflorescentiis partialibus brevibus, spicis paucis gracilibus zig-zag dispositis ferentibus; fructibus minimis (5 mill. diam.) sphaericis mucronatis, squamis nitidis profunde sulcatis et apice gibbosis.

Philippines (Vidal No. 3952).