

remota, inaequidistantia nec aggregata, elongato-lanceolata, 25-32 cent. longa, 3·4-5 cent. lata, utrinque aequaliter attenuata, acuminata, apice setoso-penicillata, supra opaca, subtus subconcolora, costulis 5 percursa, quarum medialis supra valdiuscula nuda, lateralibus minoribus et spinuloso-ciliatis, subtus nervis omnibus inconspicuis et nudis; spadicis spathae primariae elongato-tubulosae aculeolatae; spathae secundariae tubuloso-infundibuliformes inermes; spicae erecto-patulae, curvulae 4·5-6 cent. longae, floribus 10-12 distice utrinque praeditae: fructus parvi globosi 8-9 mill. diam. apiculati, perianthio pedicelliformi suffulti, squamis laxè imbricatis, semine subgloboso, superficie inaequali, albumine aequabili.

Philippines; Malanipa, (Moseley in Herb. Kew).

126. *Calamus Vidalianus* Becc. sp. n.—Fronidium rachis in parte basilari subtus convexa et nuda, longitudinaliter ad latera sulcata, aculeis vix reduncis dense armata, in facie superiore convexiuscula dense spinosa; segmenta aequidistantia, angustissime lanceolata, basi attenuata, apici acuminata, 30-32 cent. longa, 15-20 mill. lata, glabra, utrinque subconcolora, supra costulis 3 spinulosis percursa, subtus nuda, marginibus ciliato-spinulosis; spadicis ♀ spathae primariae tubuloso-cylindratae, in parte superiore aliquantum ampliatae, sparse aculeatae; spathae secundariae elongato-infundibuliformes inermes; spicae ad faucem spatharum callo insertae, patentia, arcuata, 5-7 cent. longa, floribus subdisticis utrinque paucis.—*C. horrens* (not of Bl.) Vidal.

Philippines (Vidal No. 933 in Herb. Kew).

127. *CALAMUS PISICARPUS* Bl.—*C. oblongus* β. Bl.—*C. verus* (not of Lour.) Mart. . . . . Moluccas

128. *CALAMUS ARUENSIS* Becc. . . . . Aru Islands.

129. *CALAMUS HOLLRUNGII* Becc. . . . . New Guinea.

130. *CALAMUS UNIFARIUS* H. Wendl. . . . . Borneo.

*C. UNIFARIUS* v. *PENTONG* Becc. . . . . Nicobar Islands.

131. *Calamus subinermis* H. Wendl. (nomen nudum in Herb. Kew.)—Scandens, robustus, vaginis 4 cent. diam. crases lignosis, apici saccatis, inermibus, indumento tenui cinereo-furfuraceo detergibili tectis, in ore nudis et oblique truncatis; frondes magnae cirro validissimo crebre aculeis reduncis semiverticillatis armato terminatae, petiolo brevi subtus inermi; segmentis magnis 50-60 cent. longis, 5·5-5 cent. latis, non numerosis, subaequidistantibus, elongato-lanceolatis, basi longe attenuatis, apici acuminatis, utrinque sub-