

notatis, pinnis utrinque 36—42 approximatis, aequidistantibus, lineari-lanceolatis, longissime acuminatis, horizontaliter patentibus, nervo medio nervis intramarginalibus validiore, atroviridulis; spadicibus masculis erectis, ramosis, ramis longissimis pendulis, spathis jam inter florescentiam exaridis.

Plantam femineam non vidi.

Patria: Guatemala?

Caudex diametro 9—11 lin., annulis 5—8 poll. distantibus. Frondes 5—6 ped. longae, vagina 12—16 poll., petiolo 12—15 poll., rhachi $2\frac{1}{2}$ — $2\frac{3}{4}$ ped., pinnis 11—12 poll. longis, $\frac{1}{2}$ poll. latis. Spadicum masculorum rami 12 poll. longi.

β. pinnis extimis retroflexo-patentibus.

23. Ch. elatior Mart. in Linnaea 1830 p. 205. Ejusd. Hist. III. p. 157 descr. fruct. Herm. Wendl. in Otto et Dietr. Gartenz. 1853 no. 22 p. 171. Kunth Enum. III. p. 171 no. 1. descr. fruct.

24. Ch. desmoncoides Herm. Wendl. in Otto et Dietr. Gartenz. 1853 no. 23 p. 177.

25. Ch. resinifera Herm. Wendl. in Otto et Dietr. Gartenz. 1853 no. 23 p. 179.

26. Ch. scandens Liebm. in Oversigt. 1845 p. 7 I. 9. Ejusd. in litt. ex Mart. Hist. III. p. 308 no. 16. Walp. Annal. III. p. 457 no. 7.

27. Ch. affinis Liebm. in Oversigt. 1845 p. 7 I. 10. Ejusd. in litt. ex Mart. Hist. III. p. 308 no. 17. Walp. Annal. III. p. 457 no. 8.

Euchamaedoreis verisimiliter adnumerandae sunt duae species pinnis in greges approximatis:

28. Ch. Klotzschiana sp. n. Caudice erecto, annulato; frondibus erecto-patentibus, pinnatis, vaginis cylindricis apice paululo apertis, petiolis subcylindricis, basi supra callosis, subcanaliculatis, rhachi supra acutangula, vagina, petiolo rhachique dorso fascia albida obsolete notatis, pinnis utrin-

que 15—18, ternis in greges suboppositas approximatis, suprema cujusque gregis maxima, elongato- vel oblongo-lanceolatis, falcatis, basi contractis, longe acuminatis, apicibus pendulis, summis confluentibus, opacis, nervis primariis 3, nervis secundariis utrinque 3—4.

Patria: Mexico.

Caudex diametro $1\frac{1}{2}$ poll. Frondes $4\frac{1}{2}$ — $5\frac{1}{2}$ ped. longae, vagina 10—12 poll., petiolo 10—12 poll., rhachi $2\frac{1}{2}$ —3 ped., pinnis 9—14 poll. longis, $2\frac{1}{2}$ —3 poll. latis.

Dixi hanc speciem insignem in honorem F. Klotzsch, Phil. Dr., viri praeclarissimi et scientissimi.

29. *Ch. glaucifolia* sp. n. Caudice erecto, remote annulato; frondibus erecto-patentibus, pinnatis, glaucescentibus, vaginis —, petiolis semiteretibus, supra excavato-canaliculatis, rhachi supra acutangula, petiolo rhachique dorso convexis concoloribusque, pinnis numerosis utrinque 70—80, inaequidistantibus, quaternis senisve in greges irregulares approximatis, lineari-lanceolatis, longissime acuminatis, inaequaliter patentibus.

Planta unica mascula in Hort. bot. Bruxell. colitur.

Patria: N. Granada, in pinetis prope pagum Chiapas.

Caudex circiter 12—15 ped. altus, diametro 1— $1\frac{1}{4}$ poll., annulis 9—15 poll. distantibus. Petiolus 10 poll. longus, rhachis 3 ped., pinnae 15—17 poll. longae, 3—4 lin. latae.

Species mihi non satis notae a C. C. Martio sub Euchamaedoreis enumeratae:

30. *Ch. conocarpa* Mart. in D'Orbigny Voyage dans l'Amérique méridionale Palmiers p. 6, t. 6 fig. 1, t. 16 fig. B. Ejusd. Hist. III. p. 307 no. 7.
31. *Ch. lanceolata* Kunth Enum. III. p. 172. Mart. in D'Orbigny Voyage dans l'Amérique méridionale Palmiers p. 4 t. 6