

**Chelyocarpus** Dammer gen. nov. — Truncus inermis, annulatus; folia petiolata, flabellata, petiolo inermi sectione transversali rhomboideo, marginibus obtusis, subtus et supra leviter bicanaliculato, ligula ventrali triangulata, ligula dorsali brevior, rhachide nulla, lamina usque ad basin pluripartita, segmentis elongato-cuneatis, apice laciniatis, laciniis subintegris, apice minute bidentatis; inflorescentia?; flores?; fructus apocarpus, globosus, basi perianthio (paulum aucto?) suffultus, perianthio connato 4- (an semper?) lobo, lobis coriaceis rotundatis, epicarpio tessellato pseudo-suberoso, ei fructuum *Pholidocarpi* simillimo, mesocarpio grumoso, endocarpio tenui osseo intus testa hyalina brunneo-punctata. Semen liberum, globosum rhaphe lata a basi usque ad apicem fere extensa, rhapsos ramis paucis latis parum ramosis superficiem seminis leviter cerebriformem formantibus, endospermo solido, osseo, e cellulis elongatis, incrassatis, multo perforatis, consistente, embryone parvo laterali.

**Ch. Ulei** Dammer n. sp. — Truncus annulatus, inermis, 7—8 cm diametro 2—3, rarius ad 5 cm altus. Folia petiolata, flabellata, petiolo inermi ad 75 cm longo et longiore, basi 5—7 mm lata, 3 mm crasso, apicem versus sensim attenuato, ligula ventrali triangulari 8 mm longa, 10 mm lata, ligula dorsali triangulari 2 mm longa, lamina 35—45 cm longa, ambitu semiorbiculata, supra viridia, subtus albida, rhachide nulla, usque ad basin 7—13-partita, segmentis elongato-cuneatis 35—45 cm longis, 4—10 cm latis, apice 3—5-fidis, laciniis 5—10 cm longis, apice minute bilobis, lobis 1—2 mm longis. Fructus globosus 23—25 mm diametro.

Brasilia: Prov. Amazonas: pr. fluv. Juruá Miry pr. Belem, fruct. mense Septemb. 1901 (ULE!).