

armata; spicae 6-7 cent. longae, floribus numerosis spiraliter dispositis, spathellula elongata pedicelliformi suffultis.

Borneo; Labuan (Lobb in Herb. Kew).

⊕ ⊕ Fruit not resiniferous.

40. DAEMONOROPS GRACILIPES (*Calamus* Miq.) Becc.—*Dæm.*  
*longipes* (not of Mart.) Miq.—*Calamus accedens* v.  
*brevispatha* Bl. . . . . Sumatra.

41. DAEMONOROPS LEPTOPUS (*Calamus* Griff). Mart.  
—Malayan Peninsula.

42. **Daemonorops erinaceus** Becc.—Caudices scandentes, vaginati 2·5 cent. diam.; vaginae apice conspicue saccatae ibique et in ore spiculis brunneis criniformibus dense vestitae, adjectis spinis non-nullis acicularibus longissimis (5-8 cent.), caeterum spinis elongatis angustissimis inaequalibus saepe setiformibus fuscescentibus basi in membranulis cristaeformibus aggregatis et per series appropinquatas obliquas dispositis, armatae, et indumento cincrescente indutae; frondium petiolus breviusculus (25 cent. long.) supra late canaliculatus subtus rotundatus et in medio inermis ad margines spinis validis et longis, minoribus immixtis, inordinate armatus; rachis griseofurfuracea apice cirro elongato semiverticillatim aculeato terminata; segmenta numerosa crebre aequidistantia angusta 20-30 cent. longa, 13-15 mill. lata, basi valde attenuata, apice longe acuminata, supra in costa media acuta dense spinulosa, subtus nervis 3 tenuissimis dense ciliato spinulosis percursa, et paleolis brunneis minutissimis conspersa, marginibus crebre et acute spinulosis.

Borneo; Sarawak (Beccari).

43. DAEMONOROPS OXYCARPUS Becc. . . . . Borneo.

♦ ♦ Mouth of leaf-sheaths armed with long, flat, erect spines; fruit not resiniferous.

44. **Daemonorops microstachys** Becc. sp. n.—Humilis, erectus, metralis; frondium vagina spinis complanatis subhorizontaliter seriatis armata, in petiolum longiusculum supra nudum subtus in medio aculeatum, ad margines praecipue prope basin spinis rectis 2-3 cent. longis patentibus armatum attenuata; rachis non cirrifera; segmenta satis numerosa aequidistanter remotiuscula, anguste lanceolato-ensiformia, supra costa media nuda et nervis duobus setulosis percursa, subtus tantum in costa media ciliolata; spadix gracillimus (± 20 cent. longus) pedicello gracili compresso ad margines spinoso, panicula densa inflorescentiis partialibus paucis, spathis circiter 5, quarum exterior persistens concavo-auriculaeformis elongata, extus