

toso-ciliatis, interioribus linearibus reflexis apice extus tomentosis, corollis basi extus glabris.

HAB. Woods, Organ Mountains, at an elevation of about 3500 feet. Fl. July.

Frutex diffusus, spinosus. Spinæ geminæ, 12-15 lin. longæ. Folia $3\frac{1}{2}$ - $4\frac{1}{2}$ poll. longa, 18-21 lin. lata. Pedunculi 1-4-cephali pubescenti-tomentosi, 6-9 lin. longi, basi bracteati, bracteis linear-lanceolatis, pungentibus.

5794. *Flotovia (Ernesia) leptacantha*; foliis breve petiolatis apice spinosis elliptico-oblongis utrinque obtusis supra glabris nitidis subtus ramulisque villosis, involucri squamis inermibus, exterioribus ovatis margine tomentoso-ciliatis, interioribus linearibus reflexis apice extus villosis, corollis basi extus glabris.

HAB. In ravines, near the summit of the Organ Mountains. Fl. March.

Frutex ramosissimus, spinosus 8-12-pedalis. Folia trinervia, 10-12 lin. longa, 4-4 $\frac{1}{2}$ lin. lata. Capitula subsessilia, 20-flora.

My n. 516, also from the Organ Mountains, but at a lower elevation, is a variety of this plant, with leaves nearly double the size, and less villous beneath.

5793. *Achyrophorus (Oreophila) Brasiliensis*; caule glabro striato ramoso foliis duplo breviore, foliis radicalibus longe petiolatis linear-lanceolatis subsinuato-dentatis acutiusculis supra sparse pilosis, summis linearibus integerrimis, pedunculis elongatis bracteatis unifloris, involucri subtomentoso-pubescentis squamis linear-lanceolatis, intimis longe acuminatis, acheniis glabris vix rostratis.

HAB. Moist open places, summit of the Organ Mountains. Fl. March.

Herba perennis. Caulis erectus, ramosus, 1 $\frac{1}{2}$ -pedalis. Folia radicalia 8-9 poll. longa, 4-5 lin. lata, basi in petiolum attenuata. Involucri squamæ biserialis, basi calyculatæ. Pappus uniserialis, setis filiformibus, plumosis.

LOBELIACEÆ.

5798. *Lobelia (Rapuntium) Organensis*; glaberrima, caule