

## APOGON. E.

<i>Receptaculum</i> nudum. <i>Pappus</i> 0. <i>Involucrum</i> octophyllum serie duplici.	<i>Receptacle</i> naked. <i>Pappus</i> 0. <i>Involucrum</i> 8-leaved in a double series.
--	--

## 1. HUMILIS. E.

*Root* annual? *Stem* 6—12 inches high, branching, glabrous. *Root leaves* oblong, narrow, slightly obovate, sessile. The stem *leaves* strap-shaped, acute, entire, sessile and slightly glaucous. *Flowers* terminal and somewhat umbellate, with two or more leaves sheathing the base of each umbel; perhaps 1 small leaf for each peduncle. *Peduncles* 3—8, 1—2 inches long, sometimes though rarely compound. *Involucrum* generally 8-leaved; leaves ovate, acuminate, glaucous, a little hairy and closely united at base but seeming to form two rows. *Florets* ligulate, few, (8—10) small, yellow, a little longer than the involucrum. *Receptacle* naked, flat. *Seeds* somewhat lanceolate, furrowed, transversely striate and without even the vestige of a pappus, as far at least, as the limited opportunities which I have had for examining it, have enabled me to ascertain.

Grows, though very rare, in the low and middle country of Carolina.

Found many years ago along the road between Jacksonborough and Ashepoo-Ferry. Sent to me recently from Augusta, Georgia, by Dr. Leavenworth.

Flowers April.

## STOKESIA. L'HERITIER.

<i>Receptaculum</i> nudum. <i>Pappus</i> 4-setosus. <i>Involucrum</i> foliaceum, subimbricatum. <i>Corolla</i> radiata; corollulis radii infundibuliformibus, irregularibus.	<i>Receptacle</i> naked. <i>Pappus</i> composed of 4 bristles. <i>Involucrum</i> leafy, somewhat imbricate. <i>Corolla</i> radiating; florets of the ray funnel shaped, irregular.
--	--

## 1. CYANEA.

*Root* perennial. *Stem* leafy. *Leaves* lanceolate. *Peduncles* axillary, 1-flowered. *Flowers* large, blue or purple, very handsome. Pursh.

With this plant I am entirely unacquainted.

Grows in Carolina.

Flowers.