

384. CHONDRILLA. G. pl. 1235.

Cal. calyculatus; flosculi multiplici serie; recept. nudum; sem. muricata; pappus capillaris stipite longo superne attenuato.

1. C. *junccea*, fol. radicalibus runcinatis, caulinis linearibus integris. L. Jacq. Austr. t. 427.

In calidioribus: Basileæ versus *Crenzach*, circa *Huningam* minorem, et pontem *Wiesæ* etc. *bey der Horburg*, in M. *Farnsberg*. Frequens in Valesia, circa *Branson*, *Fouly* etc. 17. Ego legi ad oras vinearum et agrorum circa *Aven* Valesiæ.

(Fol. caulina spinoso-ciliata, floscul. cœrulei.)

2. C. *Gaudini*, foliis inferioribus subovato-lanceolatis runcinatis spinosis, superioribus linearibus integris ciliatis. *Gaud.* in *Murith Guide du Botan.* p. 61.

In Valesia circa arcem *Tourbillon*.

(Videtur nova. *Gaud.*)

3. C. *prenanthoides*. Vill. C. fol. radicalibus linear-i-lanceolatis remote dentatis, caulinis linearibus integerrimis caule dichotomo, floribus fastigiatis. H.

Rarissimam hanc plantam misit am. Dr. *Custer* a Cel. *Villarsio*, postea a Cel. Dr. *Zollis*