

594. APARGIA. *Gen. pl. 1232.*

1. *A. scapis ramosis declinatis squamosis, foliis lanceolatis runcinato-pinnatifidis glabriusculis.* *Willd. sp. pl. 3. p. 1550.*

*Leontodon autumnale. Sp. pl. 1123.*

*Hedypnois autumnalis. Smith fl. brit. 2. p. 826.*

*Icon. Fl. dan. 501. Engl. bot. 830.*

In pastures and on road sides : New England. 24. Aug.

Sept. v. v. Flowers golden-yellow, outside red. It is probably an introduced plant.

595. SCORZONERA. *Gen. pl. 1230.*

1. *S. caule erecto 3-floro, pedunculis elongatis, foliis lanceolatis acutissimis acute-pinnatifidis interdum rariter dentatis.* *Mich. fl. amer. 2. p. 89.* *pinnatifida.*

*Leontodon carolinianum. Walt. fl. car. 192.*

In waste places : Carolina. *Michaux. +.*

596. LEONTODON. *Gen. pl. 1237.*

1. *L. calyce exteriore reflexo, scapo unifloro, foliis runcinatis glabris, laciniis lanceolatis dentatis.* *Willd. sp. Taraxacum. pl. 3. p. 1544.*

*Icon. Fl. dan. 574. Engl. bot. 510.*

Common in fields and grass plots ; probably introduced.

24. April—July. v. v. Flowers yellow ; it is generally known by the name of *Dandelion*. There is another species, found in shady wet woods, which I consider indigenous, but I have no materials in hand to give its description.

597. CHONDRILLA. *Gen. pl. 1235.*

1. *C. glaberrimum, multicaule ; foliis linear-lanceolatis integerrimis, pedunculis 2-3. longissimis subaphyllis unifloris.*

*lævigata.*

In Lower Virginia. ♂. Aug. Sept. v. v. Flowers bright yellow ; pappus turns a pale red ; the peduncles are from six to ten inches long, with a few linear, small leaves.