

Males distr. Hierapetra (Leon.); Zante: in colle Hagios Joannes (Mazz.).  
Apr. Jun. ☉.

**68. Lagoseris** M. B. fl. taur. cauc. III. p. 538.

1. *L. bifida* Vis. stirp. dalm. p. 19 t. 7 (*Trichocrepis*); Koch. syn. p. 435; Bois. fl. or. III. p. 881; Gelmi in bull. soc. bot. ital. 1869 p. 450; Hal. Beitr. fl. Achaia p. 25; Haussk. symb. p. 138. — *Crepis nemausensis* Sieb. avis p. 5; Ch. et B. exp. p. 234, Fl. pelop. p. 54, non Gou. — *Pterotheca bifida* Fisch. et Mey. ind. hort. petrop. 1837 n. 1736. — *P. nemausensis* Rehb. fl. exc. p. 269, Raul. cret. p. 800; non Cass. — Exsicc.: Heldr. herb. norm. n. 1263.

Hispidula; caulibus erectis vel adscendentibus, nudis, superne ramosis; foliis radicalibus, rosulatis, oblongis, in petiolum attenuatis, dentatis vel sublyrato-pinnatifidis; involucri phyllis externis calyculiformibus, internis 3—4 plo longioribus, glanduloso-pilosis vel glabratis; ligulis flavis; acheniis apice attenuatis, marginalibus laevibus, oblongis, corticosis, dorso convexis, ventre obtuse carinatis, bisulcatis, interioribus cylindricis, striatis, aliis magis externis scabridis, ceteris laevibus. — *L. nemausensi* Koch affinis, quae differt acheniis marginalibus ventre 3—5 alatis.

In collibus, saxosis, incultis regionis inferioris et montanae. Thessalia: mt. Ghavellu, ad monasterium Korona (Haussk.); Boeotia: pr. Thebas (Heldr.); Attica: pr. Athenas, mt. Lycabettus, Turcovuni, Parnes, pr. Palaeokundura ad mt. Pateras (Heldr.); Achaia: mt. Panachaicon (Hal.); Arcadia: pr. Tripolis, Alonistena, Nemnitza (Heldr.), Phigalea (Chaub.); Laconia: ad radices mt. Taygetos (Zahn); Creta (Raul.). — Mart. Jun. ☉.

16. Tribus. **HYOSERIDEAE** Less. syn. p. 127.

**69. Cichorium** L.

1. Sectio. *Eucichorium* DC. pr. VII. p. 84. — Capitula multiflora; rami inermes.

1. *C. intybus* L. sp. p. 813; S. et S. pr. II. p. 146; Pieri corc. fl. p. 115; Dallap. prosp. p. 108; Ch. et B. exp. p. 239, Fl. pelop. p. 56; Fraas fl. class. p. 197; Raul. cret. p. 795; Bois. fl. or. III. p. 715; Heldr. fl. cephal. p. 48; Hal. in z. b. G. 1888 p. 759, Beitr. Fl. Epir. p. 30, Beitr. fl. thessal. p. 15; Form. in D. bot. Mon. 1890 p. 14, in Ver. Brünn 1895 p. 25, 1896 p. 39, 1897 p. 30; Haussk. symb. p. 142. — Icon: Fl. dan. t. 907. — Exsicc.: Sint. et Bornm. it. ture. n. 1341.

Perenne, laete virens; caule erecto, aspero, divaricatim ramoso; foliis radicalibus rosulatis, runcinatis, caulinis paucis, lanceolatis, semi-amplexicaulibus; capitulis lateralibus sessilibus, geminis ternisve, termi-

viagg. Creta p. 71. — Exsicc.: Heldr. pl. gr. n. 1474, herb. norm. n. 1661; Rev. pl. cret. n. 93, in Baen. herb. europ. n. 4860; Dörfl. fl. aeg. n. 50.

Perenne, glaucovirens; caulibus nanis, a basi intricatim ramosissimis, ramis demum apice spinescentibus; foliis radicalibus rosulatis, sinuato-dentatis vel pinnatifido-lyratis, caulinis linearibus, saepius integris; capitulis in dichotomiis sessilibus, solitariis, rarius 2—3 aggregatis, terminalibus paucis, in apice ramulorum non incrassatorum insidentibus; involucri glabriusculi phyllis externis ovatis, obtusis, internis linearibus, triplo longioribus; pappo setoso-subpaleaceo, achenio 8—10 plo brevior.

In saxosis et arenosis maritimis. Attica: pr. Eleusis, Megara (Sprun.), ad Phalerum, pr. Laurion (Hausk.); in isthmo Corinthiaco pr. Kalamaki, Corinthus; insula Metopi pr. Aegina (Heldr.); Cycladum insula Syra (Weiss), Naxos (Leon.), Tenos, Mykonos (Heldr.); Laconia: a Marathonisi usque promontorium Tenare (Chaub.); Creta: pr. Canea (Tourn. voy. p. 10), Kissamos (Rev.), Khalepa, Suda, Askyphos, insula Gaidaronisi (Raul.); Cythaera: pr. Kapsali, San Nicolo (Spreitz.); Cephalonia: pr. Drapano (Heldr.). — Jun. Aug. 4.

2 × 3. *C. pumilum* × *spinosum*. — *C. hybridum* Heldr. herb. gr. norm. n. 1662. — Habitu inter parentes intermedia, a *C. pumilo* caulibus intricatim ramosissimis, ramulis demum non incrassatis, capitulis paucifloris; a *C. spinoso* radice annua, caulibus elatioribus, ramulis vix spinescentibus differt. Achenia abortiva. — In oropedio Anoraera insulae Mykonos Cycladum, inter parentes (Heldr.). —

## 70. *Hyoseris* L. gen. n. 916.

1. *H. radiata* L. sp. p. 808; Weiss in z. b. G. 1869 p 46. — Huc probabiliter: *H. lucida* Sieb. avis p. 5; Ch. et B. exp. p. 237, Fl. pelop. p. 55; vix L. mant. p. 103, quae calyculo biseriali, involuero fructifero erecto, nec stellatim expanso et foliorum segmentis imbricatis integris obtusioribus a *H. radiata* distinguitur. — Icon: Rchb. t. 9, f. 1. — Exsicc.: Dörfl. fl. aeg. n. 19.

Perennis; radice crassa verticali; foliis radicalibus, rosulatis, glabris vel ciliatis, saepe pulverulentis, in segmenta rhomboidea, dentata vel incisa runcinatim pinnatipartitis; scapis monocephalis, folia superantibus, fructiferis vix incrassatis; calyculi squamis involuero triplo brevioribus; involucri phyllis oblongo-linearibus, fructiferis stellatim expansis; acheniis marginalibus compressis, exalatis, oblongo-linearibus, striatis, pappo paleolato brevissimo, ceteris oblongo-linearibus, late alatis, costatis, margine scabris, pappo radiis numerosis, aliis setiformibus, aliis basi paleaceo-dilatatis constante superatis, intimis anguste linearibus, exalatis. — Planta graeca, saltem illa e Naxos, foliis crassiusculis, glaucovirentibus, segmentis parvis, dentatis, involucri phyllis latioribus et pappi setis basi parum paleaceo-dilatatis, magis piliformibus, ab occidentali discedit et subspeciem propriam — *graecam* — sistere videtur.