

sieca adpressa, madida erecto-patentia, e basi late vaginantia, subito lanceolata, superne subfuscata, subacuta, usque ad 6 mm longa, in parte vaginans 1 mm longa, in medio folii 0,5 mm lata, margine in toto anguste subfuscata, supra medium serrata, serratulis ca. 10–12, acutis, unicellularibus, subfuscis; cellulis vaginarum pellucidis, irregulariter quadratis, lamellae latitudinem folii totam occupantes, in sectione transversali ab uno strato cellularum 4–5 constructae, cellula marginalis paullum major, subtrotunda, papillosa. Costa lata, subcontinua, viridis, superne distinete serrata. Caetera desund.

Kiushiu: Prov. Higo, Kitamuki-Berg (Leg. N. TAKAKI Typus in Herb. K. SAKURAI 6056 4 Nov. 1934).

N.B. Ohne weiteres *Attenuata*-Gruppe. Man kann aber von Habitus und Blattbau leicht bestimmen.

Die Zeichnungen wurden alle von Herrn T. OSADA angefertigt.

Plantæ Boninenses Novæ vel Criticæ. VI.¹⁾

auctore

Takasi Tuyama.

With 4 Text-figures.

Received April 10, 1936.

SECTIONES GENERIS *Crepidiastrum*

Crepidiastrum NAKAI in Bot. Mag. Tokyo XXXIV, p. 147 (1920).

Sect. 1. *Macrocrepidiastrum* NAKAI et TUYAMA sect. nov.

Cacalia sect. *Dendrocacalia* NAKAI in Bot. Mag. Tokyo XXIX, p. 12 (1915) pro parte.

Crepidiastrum sect. *Monostemma* NAKAI in Bot. Mag. Tokyo XXXIV, p. 148 (1920), pro parte.

Frutex usque ad 1–2 m. altus simplex vel ramosus; folia integra ad apicem rami congesta; involucrum semper 5-phylum; corolla alba; pappi albi vel rufescentes.

¹⁾ Contribution from the Laboratory of Systematic Botany (Prof. T. NAKAI) of the Botanical Institute, Faculty of Science, Tokyo Imperial University.

²⁾ The expense of the study has been partly defrayed from the subsidy granted by the Foundation for the Promotion of Scientific and Industrial Research of Japan.