

especially toward the base, with some coarse salient teeth. Flowers larger than in any genuine *Stephanomeria*; the involucre fully half an inch in length, of more rigid and imbricated scales than in that genus; the ligules about three lines long, scarcely exceeding the tube of anthers. Achenia, still young, short, truncate, costate, with wide intervals, not at all narrowed at the summit. Bristles of the pappus 15 to 20, uniserial, somewhat coalescent at the base, 3 lines long, equalling the tube of the corolla, barbellate-plumose after the manner of a *Liatris*. Plant apparently as much as three feet high; the base of the stem not seen.

*CREPIS ANDERSONII*: glaber (capitulis exceptis); caudice perenni folia oblongo-spathulata vel oblongo-lanceolata laciniato-pinnatifida seu dentata subruncinata scaposque corymboso-ramosos proferente, ramis monocephalis; involucre glanduloso-pubero vel glabrato 2-3-seriali, squamis lanceolatis, interioribus attenuato-acuminatis discum fructiferum adæquantibus; acheniis fusiformibus sensim brevi-rostratis angulato-pluricostatis. — Nevada, in the vicinity of Carson, City, chiefly in low grounds. Leaves 2 to 5 inches long, including the tapering base or petiole, sometimes only denticulate or dentate, sometimes pinnatifid and with the lobes lacinate-toothed. Scapes from 10 to 24 inches high, angled and strongly striate, usually leafless and with a few linear bracts. Heads three fourths of an inch in length, or sometimes smaller; the involucre more imbricated than in *C. occidentalis* and *C. runcinata*. Flowers golden yellow. Achenia about 3 lines long, smooth, angled with salient ribs and deep furrows. Pappus soft and white.

*HIERACIUM BREWERI*: humilis (triuncialis ad subpedalem); foliis uniformibus oblongo-spathulatis integerrimis cum caule usque ad apicem folioso pilis longis mollibus albo-villosissimis; panicula corymbosa polycephala; capitulis parvis 10-15-floris; involucre subimbricato cylindræo vix glanduloso cum pedicellis parce piloso; floribus flavis; acheniis oblongo-linearibus haud apice vix basi angustatis. — Silver Valley in the Sierra Nevada, alt. 7,350 feet, and (a dwarfed alpine state) Silver Mountain, alt. 11,000 feet. Somewhat resembling the most hairy forms of *H. Scouleri*, Hook.; but dwarf, very leafy to the top, and canescent with very soft shaggy hairs, except the inflorescence; the latter of very numerous heads in a crowded panicle. Involucre 3 lines long.