

ctato, hinc veluti urceolato: Pappus sessilis, 7 — 10 setulis, rigidulis, ad lentem scabris, ultrà calycem paulò exsertis compositus: Receptaculum nudum.

H. in pratis Cyrenaicæ.

Obs. Nullo semifloscularum generi adamussim congruit: Major verò cum Hieraciis, licèt adversante calyce, quàm cum reliquis consensus.

Expl. TAB. XIII. fig. 4. a semen cum pappo; *b* semiflosculus cum semine.

CREPIS NUDIFLORA: Setuloso-hirta; foliis spathulato-oblongis, dentatis; pedunculis nudis, calyculi foliolis lineari-setaceis, patentibus; corollâ calicem paulò superante; seminibus stipitatis, transversè rugosis. **TAB. XIII. fig. 2.**

Radix simplex: Reliqua planta setulis patentibus sparsa: Caules 1. decim. circa alti, ex eâdem radice plures subæquales, erectiusculi, alii aphylli, apicem versùs nunc bifidi in pedunculos unifloros, nunc parcè dichotomi; alii parcè foliosi: Folia radicalia spathulato-oblonga, ferè sessilia, dentata, dentibus porrectis, ciliata; caulina, ubi adsunt, semiamplexicaulia, cæterum reliquis conformia; floralia ad pedunculorum dichothomiam lineari-acuminata et lineari setacea: Flos paulò ultra centim. longus: Calyculi foliola lineari-setacea, patentia, inæqualia, tertiam circa calycis longitudinem æquantia: Calyx basi paulò ampliatus, cylindricus, 7 — 8 foliolis linearibus acutis, compositus, in tubum conniventibus, nervo dorsali setulis nigricantibus patentibus hirtis, itemque pilis brevibus adpressis, lente conspicuis, sparso: Corolla lutea, calyce paulò major: Ligulæ apice profundè quadridentatæ: Semen lineari-ellipticum, transversim tuberculato-rugosum, in longum stipitem attenuatum: Pappus densis pilis simplicibus compositus: Receptaculum nudum.

H. in Cyrenaicæ pascuis.

Expl. TAB. XIII. fig. 2. a semiflosculus; *b* semen magnitudine naturali; *c* idem auctum.

CREPIS NIGRICANS: Setuloso-hispida; foliis caulinis lanceolato-amplexicaulibus, sinuato-dentatis; summis integerrimis; calyculi foliolis lineari-acutis, patentibus; calycinis dorso pilis nigris hirtis. **TAB. X. fig. 3.**

Planta setulis albis, patentibus sparsa: Caulis paniculatus, foliosus, teres, ramosus, ramis divaricatis: Folia radicalia desunt: caulina ad ramorum ortum lineari-lanceolata, amplexicaulia, sinuato dentata, dentibus porrectis, basim versùs dentato-incisa; summa, ad pedunculorum originem, lineari-acuta, sessilia, integerrima: Pedunculi ex extremis ramis nudi: Calix 1. centim. longus, foliolis 8. compositus, linearibus acutis, in cylindrum conniventibus, dorso setulis longis, nigris patentibus, duplici serie dispositis, hispidus: Calyculi foliola, calyce duplò breviora, patentia, laxa, lineari-lanceolata, pilis nigris, rigidis, hispida: Corolla calyce duplò ferè longior, lutea; ligulæ truncatæ, 4 — 5 dentatæ: Semina linearia, brevia: Pappus sessilis, densissimis pilis simplicibus compositus; Receptaculum ad lentem foveolis margine membranaceis sculptum.

H. in montibus Cyrenaicæ.

Expl. TAB. X. fig. 3. a semiflosculus cum semine; *b* semen cum pappo.

CREPIS FILIFORMIS: Foliis lineari-lanceolatis, ellipticisque, integris, vel remotè sinuato-dentatis; pedunculis aliis radicalibus, filiformibus, unifloris, nudis, aliis multifloris foliosis; bracteis linearibus in laxum calyculum confluentibus. **TAB. XXI. fig. 1.**

Radix perennis, simplex, fibrillosa: Folia radicalia primora elliptica, obsolete dentata, reliqua lineari-lanceolata, inæqualiter remotèque sinuato-dentata, omnia in petiolum attenuata, margine præsertim scabriuscula; caulina dentato-pinnatifida, summa lineari-lanceolata, integerrima: Pedunculi ex eâdem radice plures, filiformes, nudi, foliorum longitudine, uniflori: Caulis pedunculis radicalibus vix longior, uno alterove folio munitus, apicem versus florifer; bracteæ lineari-acuminatæ, ex extremis pedunculis in caliculum laxum confluentes, dorso nervoso, setuloso, hispido: Flos parvus, 7. mill. longus: Calycis foliola 9 — 10 in cylindrum conniventia, lineari-acuta, margine submembranaceo, nervoso, setuloso: Corolla lutea, calyce duplò ferè longior: Ligulæ quinque-dentatæ: Semen lineare: Pappus sessilis, densis, et scabris pilis compositus: Receptaculum nudum.

H. in Cyrenaicâ.

Expl. tab. XXI. fig. 1. a Semiflosculus cum pappo et semine; *b* calycis foliolum.

HEDYPNOIS LACINIFLORA: Foliis caulinis amplexicaulibus, lanceolatis; pedunculis elongatis, supernè incrassatis, nudis; ligulis quadrifidis, lacinulis apice glandulosis. TAB. XVII. fig. 2.

Pars inferior plantæ deest: Caulis ramosus, in pedunculos abiens filiformes, elongatos, nudos, apicem versus crassescens, hirtos, unifloros: Folia caulina ad pedunculorum basim amplexicaulia, lanceolata, utrinque hirsuta, integerrima: Flos 10 — 11 millim. longus: Calyx cylindricus, totus setulis albis, contortis, brevibus refertus; basi bracteolis nonnullis calyculatus, brevissimis, ovato-acutis, linearibusque, adpressis; foliola decem, unicâ serie disposita, lineari-acuta, dorso nervosa: Corolla flava, calycis longitudinis tertiam partem superans: Semiflosculi tubus hispidus; ligula apice 4 — 5 fida, lacinulis apice glandulâ nigrâ notatis: Semina linearia, exteriora squamis laceris, brevibus; interiora paleis quinque coronata, lineari-acuminatis: Receptaculum nudum.

H. in littore Magnæ Syrteos.

Expl. TAB. XVII. fig. 2. e Semen disci pappo paleaceo coronatum; *f* semiflosculus exterior, cum semine.

CATANANCHE LUTEA: *Linn. Sp. Pl. 1142. Desfont. Fl. atl. II. 239. Smith Fl. gr. Prodr. 145. Guss. Cat. Pl. sicil. Ten. Fl. neap. II. 176. Seb. et Mau. Fl. rom. 277. All. Fl. ped. n. 754.*

H. in Cyrenaicæ collibus.

Obs. Fig. Barr. l. c. vel pessima, vel speciem diversam refert. Fide Allionii in ligniticam civem recipio, licet in locis ab eodem memoratis eandem frustrâ perquisiverim, eiusdemque defectus in Galliâ Provinciali dubiâ confirmant.

CARDUUS TENUIFLORUS: *Willd. Sp. Pl. III. 1652. Smith Fl. gr. Prodr. II. 149. Ten. Fl. neap. Viviani Herb. Fl. lig. Decand. Fl. franç. IV. 79.*

H. in Cyrenaicæ montibus.

ATRACTYLIS CAESPITOSA: *Desfont. Fl. atl. II. 254. Tab. 225. ACARNA CAESPITOSA: Willd. Sp. Pl. III. p. III. 1700.*

H. in maritimis Tripolitanis.

STÆHLINA CHAMÆPEUCE: *Lin. Syst. Nat. ed 12: II. 538. Smith. Fl. gr. Prodr. II. p. I. 163. SERRATULA Chamæpeuce Linn. Sp. Pl. 1147.*

S. Caule fruticoso tomentoso; foliis sparsis linearibus, integerrimis, margine revolutis,




Fig. 1.




Fig. 2.



Fig. 3.

