

Tibetian Plants collected by E. Kawaguchi II

Siro KITAMURA


北村 四郎：河口慧海師採集の西藏植物 II

Crepis (Ixeridopsis) minuta KITAMURA sp. nov.

C. lactea affinis sed omnibus partibus minoribus, caulibus abbreviatis, foliis caulinis majoribus, capitulis breviter pedunculatis, corollis, achenii et pappi brevioribus.

Planta minuta 2.5cm. alta glabra. Caulis basi procumbens ramosus basibus foliorum anni praecedenti squamato. Folia inferiora rosulata spathulata 2-3cm. longa 7-8mm. lata apice rotundata integra vel mucronulata, margine irregulariter mucronulato-denticulata basi attenuata, utrinque glabra, petiolo 9-18mm. longo, folia caulina media spathulata 8-9 mm. longa 2-2.5 mm. lata. Capitula 1 mm. lata, pedunculis 5-6 mm.

longis bracteis minutis 1-1.5mm. longis ovatis instructis. Involucrum tubulosum atroviridescentia, squamae biserialales, exteriores 5, minutae inaequales ovatae vel oblongae 1-2mm. longae apice acutae vel obtuse,

Fig. 6 *Crepis minuta* KITAMURA

sae, interiores aequilongae lineares apice obtusae minute ciliolatae, hyalino-marginatae, post anthesin 8-9 mm. longae basi carinato-incrassatae. Corolla 10.8 mm. longa, ligula 1.5 mm. lata purpureo-flava, tubo 2 mm. longo. Antherae 2.5 mm. longae flavae. Achene cylindrica glabra 4.5 mm. longa 0.5 mm. lata straminea, apice truncata basi leviter contracta 10 striata, striis 5 leviter crassiusculis. Pappus albus 3.5 mm. longus, setis gracilibus apice attenuatis minut scabris. Receptaculum alveolatum nudum.

Hab. Tibet: circa Shigatse, 馬頭秘密瀧 (25 jun. 1914 E. KAWAGUCHI SM 96895 —Typus), Kabuchun rivulet (6 jul. SM 96902).

A very small *Crepis* 2-2.5 cm. high.

Crepis tibetica BABCOCK in Univ. Calif. Publ. Bot. 14: 330 (1928), The Genus *Crepis* II, 721, fig. 229 (1947).

Hab. Tibet: circa Lhasa, Panchogan (23 aug. SM 96898).

Distr. Ch'na: Szechuan, Yunnan.

Erigeron multiradiatus BENTH. in CLARKE, Comp. Ind. 56 (1876).— HOOKER f.,