

ctato, hinc veluti urceolato: Pappus sessilis, 7 — 10 setulis, rigidulis, ad lentem scabris, ultra calycem paulò exsertis compositus: Receptaculum nudum.

H. in pratis Cyrenaicæ.

*Obs.* Nullo semiflosculosarum generi adamussim congruit: Major verò cum Hieraciis, licet adversante calyce, quàm cum reliquis consensus.

*Expl. TAB. XIII. fig. 4. a* semen cum pappo; *b* semiflosculus cum semine.

**CREPIS NUDIFLORA**: Setuloso-hirta; foliis spathulato-oblongis, dentatis; pedunculis nudis, calyculi foliolis lineari-setaceis, patentibus; corollâ calicem paulò superante; seminibus stipitatis, transversè rugosis. **TAB. XIII. fig. 2.**

Radix simplex: Reliqua planta setulis patentibus sparsa: Caules 1. decim. circa alti, ex eâdem radice plures subæquales, erectiusculi, alii aphylli, apicem versùs nunc bifidi in pedunculos unifloros, nunc parcè dichotomi; alii parcè foliosi: Folia radicalia spathulato-oblonga, ferè sessilia, dentata, dentibus porrectis, ciliata; caulina, ubi adsunt, semiamplexicaulia, cæterum reliquis conformia; floralia ad pedunculorum dichothomiam lineari-acuminata et lineari setacea: Flos paulò ultra centim. longus: Calyculi foliola lineari-setacea, patentia, inæqualia, tertiam circa calycis longitudinem æquantia: Calyx basi paulò ampliatus, cylindricus, 7 — 8 foliolis linearibus acutis, compositus, in tubum conniventibus, nervo dorsali setulis nigricantibus patentibus hirtis, itemque pilis brevibus adpressis, lente conspicuis, sparso: Corolla lutea, calyce paulò major: Ligulæ apice profundè quadridentatæ: Semen lineari-ellipticum, transversim tuberculato-rugosum, in longum stipitem attenuatum: Pappus densis pilis simplicibus compositus: Receptaculum nudum.

H. in Cyrenaicæ pascuis.

*Expl. TAB. XIII. fig. 2. a* semiflosculus; *b* semen magnitudine naturali; *c* idem auctum.

**CREPIS NIGRICANS**: Setuloso-hispida; foliis caulinis lanceolato-amplexicaulibus, sinuato-dentatis; summis integerrimis; calyculi foliolis lineari-acutis, patentibus; calycinis dorso pilis nigris hirtis. **TAB. X. fig. 3.**

Planta setulis albis, patentibus sparsa: Caulis paniculatus, foliosus, teres, ramosus, ramis divaricatis: Folia radicalia desunt: caulina ad ramorum ortum lineari-lanceolata, amplexicaulia, sinuato dentata, dentibus porrectis, basim versùs dentato-incisa; summa, ad pedunculorum originem, lineari-acuta, sessilia, integerrima: Pedunculi ex extremis ramis nudi: Calix 1. centim. longus, foliolis 8. compositus, linearibus acutis, in cylindrum conniventibus, dorso setulis longis, nigris patentibus, duplici serie dispositis, hispidus: Calyculi foliola, calyce duplò breviora, patentia, laxa, lineari-lanceolata, pilis nigris, rigidis, hispida: Corolla calyce duplò ferè longior, lutea; ligulæ truncatæ, 4—5 dentatæ: Semina linearia, brevia: Pappus sessilis, densissimis pilis simplicibus compositus; Receptaculum ad lentem foveolis margine membranaceis sculptum.

H. in montibus Cyrenaicæ.

*Expl. TAB. X. fig. 3. a'* semiflosculus cum semine; *b'* semen cum pappo.

**CREPIS FILIFORMIS**: Foliis lineari-lanceolatis, ellipticisque, integris, vel remotè sinuato-dentatis; pedunculis aliis radicalibus, filiformibus, unifloris, nudis, aliis multifloris foliosis; bracteis linearibus in laxum calyculum confluentibus. **TAB. XXI. fig. 1.**



Fig. 1.<sup>a</sup>



Fig. 3.<sup>a</sup>



Fig. 2.<sup>a</sup>




Fig. 4.<sup>a</sup>