

pidato-mucronatis margine hinc inde spinuliferis sessilibus, floralibus ampliatis lato-ovatis iliciformibus venulosis margine spinulis gracilibus armatis capitula sessilia pl. m. glomerata fulcrantibus paululum superantibus; corolla albida.

**HECASTOCLEIS SHOCKLEYI.** — Very arid district, at Candelaria, Esmeralda Co., Nevada, W. S. Shockley. A remarkable addition to the few known North American *Mutisiaceæ*, to stand near *Ainsliaea*, but altogether *sui generis* and of peculiar habit. The generic name alludes to the separate enclosure of each flower in its involucre.

**CREPIS PLEUROCARPA.** Inter *C. occidentalem* et *C. acuminatam* quasi media, pube minuta cinerea demum decidua; caulibus subaphyllis; foliis runcinato-dentatis incisisve; cyma paniculiformi laxa; capitulis angustis paucifloris; acheniis oblongis sursum haud attenuatis eximie alato-10-costatis pappo paullo brevioribus. — Head-waters of the Sacramento, above Strawberry Valley, on wet slopes of the mountains, at the altitude of about 7,500 feet, 1881, Pringle. The short and thick akenes, with at least ten narrow and very salient ribs, almost wings, separated by broad grooves, distinguish this species.

**LOBELIA GATTINGERI.** *L. appendiculatæ* sat similis; floribus minoribus (lin. 3 longis); calycis lobis haud ciliatis attenuato-subulatis fere inappendiculatis basi utrinque callo minimo instructis, fructiferis capsula brevioribus; pedicellis quandoque bracteolatis. — Middle Tennessee, in springy places of calcareous bluffs and in cedar barrens, Dr. Gattinger. No. 1637 of the distribution of A. H. Curtiss, under the name of *L. leptostachys*. Flowering May and August, from a monocarpic root.

**GITHOPSIS DIFFUSA.** Demum effuse ramosissima; ramis gracilibus; foliis parvis; calycis lobis lanceolatis (basi latioribus) corollam subæquantibus ovario præsertim capsula fere linearí arcte sessili subdimidio brevioribus; seminibus turgide oblongis. — On Cucamonga Mountain, S. California, June, 1881, S. B. & W. F. Parish. The capsule opens apically, as in the original species.\* The blue corolla is only 2 lines in length.

**ANDROSACE ARIZONICA.** E grege *A. occidentalis*, tenella; scapis debilibus decumbentibus radiisque umbellæ paucifloræ capillaribus elongatis; foliis phyllisque involueri consimilibus brevibus; calycis lobis foliaceis, fructiferis accrescentibus ovatis radiato-patentibus tubo

---

\* Baillon's statement to the contrary is founded on a misapprehension, he evidently having taken a Texan *Specularia* for *Githopsis*. See Bull. Soc. Linn. Par. 304.