

Smooth Crepis.

Nat. of Britain.

Fl. June—September.

H. ○.

7. *C. foliis runcinato-pinnatifidis scabris; basi superne biennis, dentatis, calycibus muricatis. Sp. pl. 1136.*

Biennial Crepis.

Nat. of England.

Fl. July and August.

H. ♂.

8. *C. foliis radicalibus runcinatis; caulibus hastatis, calycibus subtomentosis. Sp. pl. 1133. Dioscoridis.*

Dioscorides's Crepis.

Nat. of France.

Introd. 1772, by Monf. Richard.

Fl. June and July.

H. ○.

9. *C. foliis runcinato-dentatis subincanis, pedunculis nudis unifloris, squamis calycinis marginæ albicantibus. De Lamarck encycl. 2. p. 179.*

*Crepis albida. Jacqu. ic. collect. 1. p. 81. Allion. pedem. 1. p. 219. t. 32. f. 3.*

Pale-flower'd Crepis.

Nat. of France and Italy.

Fl. July—October.

H. ♀.

10. *C. foliis oblongis duplicato-ferratis setosis, caule nudo ramoso, floribus paniculatis, calycibus cylindraceis glabris, pappo sessili. rigens.*

Brittly-leav'd Crepis.

Nat. of the Azores. Mr. Francis Masson.

Introd. 1778.

Fl. July and August.

G. H. ♀.

H. C.