

- Idianthes* Desv., Fl. Anj. 199. 1827.
Intybellia Monn., Ess. Hierac. 78. 1829.
Intybus Fries, Novit. Fl. Suec. ed. 2, 244. 1842.
Lagoseris Hoffmegg. et Link, Fl. Port. 2:149. 1820.
Lagoseris M. B., Fl. Taur. Cauc. 3:538. 1819.
Lepicaune Lapeyr., Hist. Abr. Pl. Pyr. 478. 1813.
Lepidoseris Rehb., Fl. Exc. 1:256. 1830-1832.
Limnocrepis Fourr., Ann. Soc. Linn. Lyon, n.s. 17:104. 1869.
Limnoseris Peterm., Fl. Lips. 589. 1838.
Myoseris Link, Enum. Hort. Berol. 2:291. 1822.
Nemauchenis Cass., Bull. Soc. Philom. 1818:77.
Omalocline Cass., Dict. 48:431. 1827.
Pachylepis Less., Syn. Comp. 139. 1832.
Paleyia Cass., Dict. 39:393. 1826.
Phaecasium Cass., Dict. 39:787. 1826.
Phalacroderis DC., Prod. 7:97. 1838.
Psammoseris Boiss., Diagn. ser. 1, 11:52. 1849.
Psilochaenia Nutt., Trans. Am. Phil. Soc. n.s. 7:437. 1841.
Pterotheca Cass., Bull. Soc. Philom. 1816:200.
Rhynchopappus Dulac, Fl. Hautes-Pyr. 493. 1867.
Rodigia Spr., Neu. Entd., 1:275. 1820.
Sclerolepis Monn., Ess. Hierac. 81. 1829.
Sclerophyllum Gaud., Fl. Helv. 47. 1829.
Soyeria Monn., Ess. Hierac. 75. 1829.
Succisocrepis Fourr., Ann. Soc. Linn. Lyon, n.s., 17:104, 1869.
Trichocrepis Vis., Stirp. Dalm. 18. t. 7. 1826.
Trichoseris Vis., *op. cit.* 19.
Wibelia Gaertn., May. et Scherb., Fl. Wett. 3:97. 1801.
Zacintha (Tourn.) L., Syst. ed. 1. 1735.

ORIGINAL DESCRIPTION

621. **Crepis**¹ Hieracioïdes Vaill. A. G. 1721. 47. 52. 13. 17. 18.

- CAL: *Communis* auctus, oblongus: *Squamis multis* linearibus, aequalibus, parallelis. *S. paucis* brevibus, basi incumbentibus.
 COR: *Composita* imbricata, uniformis: *corollulis hermaphroditis* plurimis, aequalibus. *Propria* monopetala, ligulata, linearis, truncata, quinquentata.
 STAM: *Filamenta* quinque, capillaria, brevissima. *Anthera* cylindracea, subulata.
 PIST: *Germen* infra corollam propriam. *Stylus* filiformis, longitudine staminum. *Stigmata* duo, reflexa.
 PER: nullum. *Calyx* subrotundus.
 SEM: solitaria, oblonga, coronata *Pappo* longo, plumoso.
 REC: nudum.

EXPANDED DESCRIPTION

Perennial, biennial, or annual herbs; plant \pm pubescent or glabrous; rhizomatous or with a deeply penetrating, woody or ephemeral root; caudex \pm woody, rarely suffruticulate, leafy; caudical leaves usually rosulate, many or few, persistent or ephemeral; cauline leaves numerous, few or none, the lower similar to the caudical ones or (sec. *Pyrimachos*) cataphyllous, the others gradually or abruptly reduced, sometimes all or mostly bractlike; stem or stems erect or decumbent, \pm branched or

¹ *Crepis* Vaill. is referred to *Sonchus* Tourn. by Linnaeus (Gen. ed. 1. 240. no. 617. 1737).

Subsection C.—**Corymbiformae** subsec. nov. Planta 2–12 dm alta; folia caudicalia ampla elliptica vel oblanceolata; folia caulina plerumque numerosa conspicua lanceolata vel elliptica; caulis robustus paniculatus; inflorescentia corymbiformis vel racemiformis; capitula magna vel mediocria 12–100-flora; involucria setosa vel pubescentia vel tomentosa.

Plant 2–12 (mostly 3–6) dm high; caudical leaves broad, elliptic or oblanceolate; cauline leaves usually numerous, gradually reduced, conspicuous, lanceolate or elliptic; stems robust, paniculately branched; inflorescence corymbiform or racemiform; heads large or medium, 12–100-flowered; involucria setose, pubescent with or without glands, or tomentose. Type species: *C. pannonica* (Jacq.) Koch.

Subsection D.—**Subcorymbiformae** subsec. nov. Planta 1–4.5 dm alta; folia caudicalia ampla oblanceolata; folia caulina plerumque parva bracteiformia; caulis rectus robustiusculus vel tenuis furcatus; inflorescentia subcorymbiformis; capitula mediocria vel magna 50–80-flora; involucria tomentosa saepe pubescentia.

Plant 1–4.5 dm high; caudical leaves broad, oblanceolate; cauline leaves mostly small, bractlike; stem erect, rather stout or slender, 1–3-furcate; inflorescence subcorymbiform; heads medium or large, 50–80-flowered; involucria \pm tomentose, often pubescent. Type species: *C. auriculaefolia* Sieber.

Subsection E.—**Divaricatae** subsec. nov. Planta 0.3–5 dm alta; folia caudicalia ampla oblanceolata; folia caulina plerumque parva bracteiformia; caulis tenuis furcatus, ramis paucis divaricatis; inflorescentia diffusa; capitula mediocria 15–80-flora; involucria tomentosa interdum pubescentia vel setulosa.

Plant 0.3–5 (mostly 1–3.5) dm high; caudical leaves broad, oblanceolate; cauline leaves mostly small, bractlike; stem slender, furcate, divaricately few-branched; inflorescence diffusely cymose; heads medium, 15–80-flowered; involucria tomentose, sometimes pubescent or setulose. Type species: *C. Raulini* Boiss.

Subsection F.—**Strictae** subsec. nov. Planta 1.5–4.5 dm alta; folia caudicalia linearia vel anguste oblanceolata; folia caulina similia gradatim reducta; caules tenui ramosi, ramis paucis elongatis strictis; inflorescentia cymosa; capitula magna mediocria vel parva 11–40-flora; involucria tomentosa vel pubescentia.

Plant 1.5–4.5 dm high; caudical leaves linear or narrowly oblanceolate; cauline leaves similar, gradually reduced; stems slender, few-branched, the branches elongated, strict; inflorescence cymose; heads large, medium, or small, 11–40-flowered; involucria tomentose or pubescent. Type species: *C. turcomanica* H. Krasch.

Section 11.—**Macropodes** sec. nov. Herbae perennes; planta pubescens; radix robusta lignea elongata; caudex ligneus simplex vel furcatus foliatus; folia infera oblanceolata dentata vel pinnatifida petiolata; folia superna bracteiformia; caules 0.2–3.5 dm alti tenui scapiformes vel 1-furcati cum 2–4 capitulis; capitula mediocria vel parva 11–60-flora; corolla 10–18 mm longa, tubo 2.5–6.5 mm longo; achaenia straminea fusca vel nigra 4–9.5 mm longa 10–20-costata vel -striata; pappus albus flavidulus vel fuscidulus 3–8 mm longus.

Perennial herbs; plant pubescent; root stout, woody, elongated into a taproot; caudex leafy, 1- or several-stemmed; lower leaves oblanceolate, denticulate, dentate, pinnatifid or bipinnate, petiolate; upper leaves all or mostly bractlike; stem or stems slender, scapiform, 1-headed, or sometimes furcate and 2- or rarely 3–4-headed; heads medium or small, with 11–60 or more florets; longest outer bracts $\frac{1}{4}$ – $\frac{2}{3}$ (mostly $\frac{1}{3}$ – $\frac{1}{2}$) as long as the inner; inner bracts becoming convex and indurate or slightly thickened but otherwise little changed at maturity, or becoming definitely carinate and spongy-thickened dorsally, glabrous or pubescent on inner face; corolla 10–18 mm long, the tube 2.5–6.5 mm long; achenes ranging in color from stramineous or pale brown to dark brown, dark purple, or black, 4–9.5 mm long, fusiform or columnar, sometimes strongly attenuate or rarely shortly beaked, 10–20-striate or -ribbed, the striae or ribs unequal or equal; pappus white, yellowish, or tawny, 3–8 mm long. Macropodes, from *makros*, long, + *pous*, *podos*, a foot; referring to the long taproot. Type species: *C. oreades* Schrenk.

Section 12.—**Ixeridopsis** sec. nov. Herbae perennes; planta glabra; radix tenuis lignea elongata; caudex tenuis ligneus simplex vel furcatus interdum elongatus foliatus; folia infera ovata elliptica vel oblanceolata sessilia vel petiolata integra denticulata dentata vel subpinnatifida; folia superna similia vel bracteiformia; caules 0.2–3.5 dm alti tenui ramosi; inflorescentia corymbiformis vel diffusa; capitula mediocria vel parva 6–13-flora; involucrum cylindricum tenue glabrum, squamis exterioribus 5–8 ovatis brevissimis, interioribus 8–10 lanceolatis 3–7-plo longioribus; receptaculum nudum; corolla 7–14 mm longa, tubo 2.5–5 mm longo; achaenia straminea vel fusca 4–8 mm longa teretia ad summitatem paululum attenuata vel bressime rostrata 10-costata vel -striata; pappus albus vel flavidus 4–6.5 mm longus.

Perennial herbs; plant glabrous; root slender, woody, elongated into a taproot; caudex slender, woody, simple or furcate, sometimes much elongated and branched, leafy; lower leaves ovate,