

in Australia, New Zealand, and in North America, where, however, it may have been introduced.

Crepis japonica, *Benth.*; *Fl. Brit. Ind.* iii. p. 395.—Shan hills plateau at 5000 feet.

Afghanistan eastward to Japan and southward to Australia; also in Mauritius and South Africa, where it may have been introduced.

Crepis acaulis, *Hook. f. in Fl. Brit. Ind.* iii. p. 396.—Shan hills at 3000 feet.

Very widely spread in India.

Crepis (§ Youngia) subscaposa, *Coll. et Hemsl.*, n. sp.

Herba biennis? hispidula, pilis simplicibus, 9–18 poll. alta, caulibus aphyllis gracilibus superne pauciramosis. *Folia* rosulata, crassiuscula, obovato-oblonga vel spathulata, in petiolum brevem attenuata, 2–3 poll. longa, apice rotundata, pauci callosodenticulata, utrinque hispidula. *Capitula* pauca, circiter 5–7, corymbosa, longe pedunculata, 6–8 lineas diametro, circiter 30-flora; involucri calyculati bracteæ hispidulæ, intimæ uniseriatæ, lineari-oblongæ, obtusæ, marginibus scariosæ, pappum triente breviores; receptaculum parvum, nudum; ligulæ breves, brevissime calloso 5-dentatæ, extus infra medium puberulæ; antheræ basi breviter setaceo-caudatæ. *Achænia* recta, subcompressa, gracilia, pluricostata, nuda, supra medium hispidula, sursum attenuata, pappo longiora; pappi setæ albæ, molles, uniseriatæ, simplices, in annulum deciduæ.

Shan hills at 3000 to 5000 feet.

Not closely allied to any species, and in habit resembling some of the small-headed *Hieracia*.

Crepis (§ Youngia) chloroclada, *Coll. et Hemsl.*, n. sp.

Herba perennis, 12–20 poll. alta, a basi multiramosa, fere undique glabra, caulibus ramisque gracilibus rigidis lignescensibus angulatis viridibus. *Folia* radicalia non visa, caulina coriacea, glabra, linearia, 1–1½ poll. longa, vel interdum ad bracteas breves subulatas reducta. *Capitula* angusta, 7–10-flora, longe graciliterque pedunculata; involucri calyculati bracteæ circiter 7, lineari-oblongæ, obtusæ, dorso hispidulæ;