

708. *MICRANTHA*. Capitula α -flora. Involvari bracteæ parvæ. Achenia exteriora paleis induratis acris lobata pappi setis brevibus subplumosa, interiora angusta, nuda calva raris. Herba annua, stans, cauligera, foliis pinnatifida. Capitula longè pedunculata.—*Chili*.

709. *PANICULARIA*. Capitula α -flora. Involvari bracteæ sub-2-seriatæ. Receptaculum nudum. Achenia nuda. Herba annua, erecta, glabra, foliis obtusa integris lobatis. Capitula parva, longè gracilliterque pedunculata.—*And. austr. subæqv.*

710. *CASALUARPIA*. Capitula α -flora. Involvari bracteæ sub-2-seriatæ. Receptaculum nudum. Achenia tuberculata ovata sublae apice truncato mucronata. Herba perennis, subscandens, lanuginosa, foliis radiatibus ovatis, ovale apiculatis longè 2-3-lobatis.—*Brazilia*.

Tribo XIII. Cichoriaceæ. Fus. (Lipsifera, DC.). Capitula homogena, floribus omnibus γ fertilibus v. his inde paucis sterilibus. Corolla tubo tenui, limbo in laciniam seu ligulam palmatam 5 membra, apice truncato 5-dentato expanso, sepalinis cæcis ad apicem tubi plus minus pilosis. Antheræ apice appendiculatæ, basi sagittata, auriculis antherarum distinctis mucronatis acutis v. breviter obtuso-angulatis (rursum ovula lateris appendiculatis?). Styli rami tenues, obtusiusculi v. subacuti, subvix papillosi, rursus abbreviati v. leviter dilatati. Herba, raris frutescens v. (in gen. 2 involucris) aëreæ, sæpiè simpliciter lactes, internodiis v. raris raris spinosissimis, v. (in *Stolypis*) herbæ mollioribus. Folia alterna v. radiata. Involvari bracteæ apice membranaceo-lanceolatis, subequaliter 1-2-seriatæ v. calyculatis, v. perianthiatis raris multimeriatis lobatis. Corollæ apicis flava, interiorum nervibus v. raris, raris albis.

Tribo naturalissima, quod forma nuda aëreæ, difficillime in subtribus dividitur et inter sequentes numeralis male limitata.

Subtribus 1. *Stolypis*. Herba aëreæ, foliis involvariisque spinosissimis. Achenia a disco compressa, paleis receptaculi abbreviatis subulatis acris induratis, in alata et cum his densis; pappus 6 v. paucioribus.

711. *Stolypis*. Genus unicum.—*Sp. Mediceæ*.

Subtribus 2. *Dactyloctenium*. Aëreæ. Involvari bracteæ α -seriatæ, induratis, basi plus minus mucronatis. Achenia a disco compressa, sub-3-gona v. 2-3-angula.

712. *Dactyloctenium*. Capitula pedunculata. Receptaculum nudum v. sublobiferum. Pappus α -seriatis aëreæ.—*Ind. Javae Formosæ*.

713. *Pycnia*. Capitula magna, solitaria. Receptaculum palmatum. Achenia plana, setis v. setolis 2 longis villosis mucronatis.—*Ind. Tabaci et Otahiti*.

Subtribus 3. *Hymenocleis*. Herba. Involvari raris. Achenia apice truncata, pappo plus minus palmato v. palmato parvo cum setis alternatis mucronatis v. nudo.

Keyes: Pappus setosus, setis involucris subpalmatis in *Hymenocleis*. Achenia subtrilobata in *Hymenocleis*.

* Involvari bracteæ α -seriatæ, induratis, productis mucronatis.

714. *Catananthe*. Involvari bracteæ ovale mucronato-appendiculatæ. Receptaculum mucronatum. Pappi palmæ 5-7. Herba, foliis plerisque radiatibus linearibus integris v. pinnatifidis. Capitula longè pedunculata. Corollæ nervibus v. flava.—*Sp. Mediceæ*.

715. *Hymenocleis*. Involvari bracteæ marginè submembranaceis. Receptaculum breviter mucronatum. Pappi setis 2-3-seriatis, lobulatis, interioribus subpalmatis. Herba, foliis plerisque radiatibus dentatis v. pinnatifidis. Capitula longè pedunculata. Corollæ flava.—*Sp. Mediceæ*.

716. *Hymenocleis*. Involvari bracteæ marginè submembranaceis. Receptaculum palmatum. Pappi palmæ 3-6. Herba, foliis radiatibus dentatis v. pinnatifidis. Capitula in seipso sub-aphylla solitaria. Corollæ flava.—*Hispænia*.

** Involvari bracteæ interioribus 1-2-seriatæ, ovatis, subequalibus, exterioribus brevibus v. laevibus v. 6. Receptaculum nudum v. sublobiferum.

717. *Cichorium*. Involvari bracteæ interioribus mucronatis, disco mucronato. Pappi palmæ breves, mucronatis. Herba, raris rigidis v. spinosissimis. Capitula sessilia v. rigidè pedunculata. Corollæ corollæ.—*Hispænia, her. orbis interio, in alia loca impatiens*.