

teriora longiora, lanceolata, vel longe elliptica, 9 mm longa, 1.5 mm lata, apice obtusa; omnia extus pallide rosea, glabra. Flosculi ligulati, albi, 4 in numero. Achaenia immatura. Pappi setae albae, scabridae.

Sichuan (四川): Emei Mt. (峨眉山), in sylva, alt 2500 m, 1935-08-29, T. H. Tu (杜大华) 693 (Typus, PE).

17 南川花佩 新种 图 7

Faberia nanchuanensis Shih, sp. nov.

Ex affinitate *F. lanceifoliae* Anth., ab ea differt forma foliorum.

Herba perennis. Caulis solitarius, erectus, 34 cm altus, basi 2.5 mm in diametro, infra medium paniculato-ramosus, omnino glaber. Folia basalia numerosa, longe elliptica, 8—10 cm longa, 1—1.5 cm lata, apice breviter acuminata, basi in petiolos longos cuneato-angustata, petiolis 5—8 cm longis, margine sparsim obscure repando-serrulatis; folia caulina 6 in numero, apice acuminata vel longe acuminata, longe elliptica vel ea suprema longe linearia, ea infima petiolata, petiolis 6 cm longis, ea suprema sessilia, basi cuneato-angustata; omnia textura coriacea, glabra, laevia, subtus purpurea. Calathidia 12 in numero, paniculato-disposita. Involucrum in statu explanato late cuneatum, cuius phyllis 4-seriatis, obscure imbri-catis, extremis externisque ovato-triangularibus vel longe ovatis, usque ad 5 mm longis, 1.5 mm latis, intimis internisque longe ellipticis, usque ad 1.3 cm longis, 1.5 mm latis; omnibus extus glabris, viridibus, apice acutis vel obtusis. Flosculi numerosi caerulei, ligulati. Achaenia immatura.

Sichuan (四川): Nanchuan (南川), prope fossas in locis humidis, 1957-06-28, J. H. Xiong et Z. L. Zhou (熊济华与周子林) 91698 (Typus, PE).

18 耳叶细离草 新种

Stenoseris auriculiformis Shih, sp. nov.

Species insignis foliis auriculiformi- vel codato-amplexicaulibus, a speciebus ceteris sinensibus bene distincta.

Herba annua, radicibus fibrosis numerosis. Caulis erectus, solitarius, 75 cm altus, basi 2 mm in diametro, superne corymboso-ramosus, ramis gracilibus, omnino glaber. Folia basalia sub anthesi emarcida; ea media petiolata, petiolis 3—9 cm longis, basi eorum auriculiformi-amplexicaulibus, lyrato-pinnatisecta vel -pinnatipartita, lobis lateralibus 1—2 jugis, ellipticis vel longe ellipticis, 0.7—2.5 cm longis, 0.8—1.8 cm latis, apice obtusis vel rotundatis, mucronulatis, margine integris vel paucō-serrulatis, -apiculatis, lobis terminalibus triangularibus vel cordatis, 3—4 cm longis, 2.5—3 cm latis, apice acutis, mucronulatis, margine repandis, paucō-serrulatis vel -apiculatis; ea media superioraque longe cordata vel -ovata, sessilia, basi valde cordato-amplexicaulia, margine integra, apice longe acuta; omnia textura tenuia, utrinque glabra. Calathidia numerosa, corymboso-disposita. Flosculi purpurei, 5 in numero, ligulatis. Involucrum anguste cylindricum, 8 mm longum, 2 mm in diametro, cuius phyllis 2-seriatis, externis brevissimis, longe ovatis, 1.8 mm longis, 0.8



图 7 南川花佩。(蔡淑琴绘)
Fig. 7 *Faberia nanchuanensis* Shih.