

Helminthotheca echioides (L.) Holub, *Folia Geobot. Phytotax.* **8**: 176 (1973).

var. **mollis** (Duby) P. D. Sell comb. nov.

Helminthia echioides var. *mollis* Duby, *Bot. Gall.* **1**: 300 (1828).

var. **pratensis** (Chevall.) P. D. Sell comb. nov.

Helminthia echioides var. *pratensis* Chevall., *Fl. Gén. Env. Par.* **2**: 352 (1827).

var. **incisa** P. D. Sell var. nov.

Holotype: Garden of 54, Cambridge Road, Impington, Cambs, v.c. 29, 52/436619, 15 July 1998, P. D. Sell no. 98/92 (**CGE**) (1 plant on 6 sheets).

Planta erecta, pilis longioribus rigidis aequa cum brevibus vestita. *Folia* magna, costam versus ad partem dimidiam lobis latis dentatis incisa. *Involuci bracteae exteriore*s ad basin cordatae.

Sonchus oleraceus L., *Sp. Pl.* 794 (1753).

var. **litoralis** P. D. Sell var. nov.

Holotype: Shingle beach between Cley and Salthouse, E. Norfolk, v.c. 27, 63/059451, 21 September 1967, P. D. Sell no. 67/1775 and N. Marchant (**CGE**).

Caulis 10–50 cm. *Folia* saepe rubripurpureotincta. *Capitula* pauca, in fasciculo denso congesta.

Sonchus asper (L.) Hill, *Herb. Brit.* **1**: 47 (1769).

var. **sabulosus** P. D. Sell var. nov.

Holotype: Stabilised shingle at back of beach, north of Dunwich, E. Suffolk, v.c. 25, 62/479709, 26 June 1990, P. D. Sell no. 90/157 and J. G. Murrell (**CGE**).

Caulis ad 50 cm. *Folia* plerumque divisa, aculeatissima. *Capitula* pauca, in fasciculo denso congesta.

Lactuca virosa L., *Sp. Pl.* 795 (1753).

forma **lactucarii** (Lamotte) P. D. Sell stat. nov.

L. lactucarii Lamotte, *Prodr. Fl. Plat. Centr.* **2**: 467 (1881).

L. scariola forme *virosa* var. *lactucarii* (Lamotte) Rouy, *Fl. France* **9**: 199 (1905).

Crepis vesicaria L., *Sp. Pl.* 805 (1753).

var. **rutilans** (Lacaita) P. D. Sell comb. et stat. nov.

C. rutilans Lacaita, *Jour. Linn. Soc. Bot.* **44**: 128 (1918).

Pilosella × *stoloniflora* (Waldst. & Kit.) F. W. Sch. & Sch. Bip. in *Flora (Regensb.)* **45**: 423 (1862).

nothosubsp. **schuriana** (Nägeli & Peter) P. D. Sell comb. nov.

Hieracium × *stoloniflorum* subsp. *schurianum* Nägeli & Peter, *Hier. Mittel-Europas: Piloselloiden* 326 (1885).

Probabiliter = **P. aurantiaca** subsp. **carpathicola** × **officinarum** subsp. **euronota**

Hieracium L.

In Section *Sabauda* (Fr.) Arv.-Touv.

Hieracium vagum Jord., *Cat. Grenoble* 21 (1849).

forma **croceostylum** (Pugsley) P. D. Sell comb. et stat. nov.

H. croceostylum Pugsley in *Jour. Linn. Soc. London (Bot.)* **54**: 310 (1948).

Hieracium prominentidens P. D. Sell nom. et stat. nov.

H. dispalatum var. *macrodon* Sudre, *Hier. Centr. France* 24 (1902), non *H. macrodon* Nägeli & Peter, *Hier. Mittel-Europas: Archieraciens* 84 (1886).

Hieracium sabaudum L., *Sp. Pl.* 804 (1753).

forma **bladonii** (Pugsley) P. D. Sell comb. et stat. nov.

H. bladonii Pugsley in *Jour. Linn. Soc. London (Bot.)* **54**: 299 (1948).

In Section *Hieracioides* Dumort.

Hieracium umbellatum L., *Sp. Pl.* 804 (1753).

var. **sarniense** P. D. Sell var. nov.

Holotype: Granite cliffs, north coast of Jersey, Channel Islands, 17 July 1901, L. V. Lester-Garland as E. F. & W. R. Linton, *Set of British Hieracia* no. 185 (**CGE**).

Caulis 10–30 cm, robustus, succulentus. *Folia* crassa, succulenta, anguste vel late elliptica, ad apicem obtusa, inferiora ad c. 5 cm lata, prope caulis basin fasciculata, sursum minora.

var. **latifolium** (W. R. Linton) P. D. Sell stat. nov.

H. umbellatum forma *latifolium* W. R. Linton, *Brit. Hier.* 98 (1905).

In Section *Foliosa* (Fr.) Arv.-Touv.

Hieracium reticulatiforme P. D. Sell nom. et stat. nov.

H. reticulatum var. *angustifolium* Pugsley in *Jour. Linn. Soc. London (Bot.)* **54**: 280 (1948), non *H. angustifolium* Hoppe, *Bot. Taschenb.* **1799**: 129 (1799).

Hieracium reayense (Pugsley) P. D. Sell stat. nov.

H. reticulatum var. *reayense* Pugsley in *Jour. Linn. Soc. London (Bot.)* **54**: 280 (1948).

Hieracium melanoglochin (E. F. Linton) P. D. Sell stat. nov.

H. corymbosum var. *melanoglochin* E. F. Linton in W. R. Linton, *Brit. Hier.* 88 (1905).

In Section *Tridentata* (Fr.) Arv.-Touv.

Hieracium charitodon P. D. Sell sp. nov.

Holotype: Waterfall at the head of Glyn Tarell, Brecon, v.c. 42, 25 July 1900, A. Ley (**CGE**).

Herba perennis aphyllopoda vel hypophyllopoda, caudice ramoso. *Caulis* 35–100 cm, pallide luteoviridis, ad basin saepe purpurascens, gracilis vel robustus, infra pilis simplicibus eglandulosis densis longis pallidis, sursum paucioribus brevioribusque, et in parte superiore pilis stellatis numerosis vestitus. *Folia* in pagina superiore pallide vel mediocriter viridia, in pagina inferiore pallidiora, 5–15, sursum gradatim decrescentia; folia basalia et caulina infera lamina 6–11 × 1.5–4.0 cm, anguste elliptica vel oblangoelliptica vel lanceolatooblonga, ad apicem obtusumucronatum vel acutum gradatim angustata, dentibus mammiformibus vel longiusculis angustisque dentata, ad basin in petiolum alatum ad 4 cm longum attenuata; caulina mediana lamina 4–9 × 1.5–4.0 cm, linearieelliptica