

amplexicaulia plus minus remote denticulata, anthodii squamæ lanceolatae apice pinnatifido-spinulosæ; involucri foliola latiora sub-spinosa, angustiora et evidentius spinosa in var: *a* ex *Pyrenæis* (*H. spinosa* DC. syn: p. 265.).

CLXX. HYPOCHÆRIS Lin: gen: ed: Schreb: n. 1246. = Spr: syst: n. 2898. = Lam: ill: t. 656. = *Achyrophoris* spec: Scop:

» *Anthod:* imbricatum. *Recept:* paleaceum. *Pappus* » radii sessilis disci stipitatus (non semper). »

Obs: Genus constans 10-12. stirpibus herbaceis fere omnibus europæis, perpaucis africanis vel americanis; caule simplici vel ramoso plus minus folioso, foliis oblongis sinuato-dentatis, anthodii squamis et pappo variis. Species pappis omnibus stipitatis constituunt gentis *Achyrophori* Scop:

• 1. *H. radicata* = L. h. Cliff: 386. = Spr: syst: III. 668. = Fl: dan: t. 150. Ic: taur: vol. 8. t. 64. et vol. 11. t. 47. (*Eur:* cultiore. *Pedem:* abunde in pratis non pinguibus et pascuis, in agro *Nicæens:* secus arva frequens All: n. 847. RE fl: tor: II. p. 53. BIROLI fl: acon: II. p. 60. Passim in pascuis et speciatim in apricis sterilibusque *di Condove* in regione dicta *la Roccea* RE fl: segus: p. 66. In *Sardin:* arvis MORIS st: sard: I. p. 28.). Perenn: Caulis subramosus nudiusculus glaber, folia scabra, pedunculi elongati superne incrassati, anthodii squamæ carina ciliatæ, flores magni pro genere, pappi omnes stipitati.

Hypochaerides dubiæ vel novæ.

7. *H. hirta* = Cyr: sec: Steud: Habui ex hb: BIROLI sub hoc nomine, nec illam video enumeratam ab Auctoribus nisi a STEUD: Planta pumila glabra, folia lineari-lanceolata denticulata, caules ramosi sub-aphylli squamosi, anthodii squamæ lineari-lanceolatæ, pappus generis. Nomen *hirtæ* non convenit plantæ meæ undique glabræ. = 8. *H. taraxacifolia* = Loisl: nouv: notic: p. 33. (*Corsica. Sardin:* in pascuis montanis MORIS st: stard: II. p. 26.). Cæspitosa glabra ramosa, radix crassa fere tuberosa, folia radicalia lineari-lanceolata longissime petiolata runcinato-pinnatifida caulina linearia integra, pedunculi squamulosi sub-æquales 1-flori, anthodii squamæ lineari-lanceolatæ margine albo - scariosæ, pappi omnes stipitati. Possideo specim: simillima ex hb: BIROLI nomine *Apargiæ taraxacoideæ* sine alia indicatione.

9. *H. Berteri* = Nob: *H....?* Berter: ined: (*Chili* in pratis arenosis). « *H. caule erecto ramoso folioso* » pilosiusculo, foliis oblongis acutis sub-sinuato-dentatis glabris costa media utrinque pilis simplicibus » hispidula, pedunculis apice sub-incrassatis 1-floris, » anthodii squamis lanceolatis hispidis margine albo-scariosis. » NOB: Culta in h. Ripul: e seminibus missis a BERTERO sine nomine specifico; an eadem ac *H. arenaria* GAUDICH: fl: Os Malouin: et SPR: c. p. 305.? Ex diagnosi Auctorum stirps nostra differt præsertim foliis vix sinuato - dentatis nec pinnatifidis. Cons: COLLA pl: rar: chil: fasc: II. p. 40. t. XXXIII.