

**Hypochoeris radicata** L. Sp. pl. 811. Var. *heterocarpha* Moris. Fl. Sard. Mm. Treskavec et Selca pl. in M.

**H. macedonica** Form. 1898. Perennis, caule subtenuē, angulato-striato, 25 cm. alto, bifolio, valde parce et sparsim setaceo, foliis radicalibus rosulatis, longe oblongo-lanceolatis, 15 cm. longis, remote repando-paucidentatis, in petiolum alatum attenuatis, subtus ad nervum medium setis brevibus obsitis et ad margines ciliatis, caulinis lineari-setaceis, capitulis mediocribus, involucri phyllis imbricatis, dorso setosis, externis lineari-setaceis vel linearibus, internis lineari-lanceolatis, longe acuminatis albidis, dorso nervo viridi lato percussis, margine parce ciliatis, flosculis marginalibus involucrum superantibus, ligulis luteis supra tubum dense-, ad dorsum valde parce hirsutis, apice profunde 5 dentatis, acheniis plano compressis scabridis, sat longe attenuato rostratis, pappo plumoso. Habitat, in reg. superiore m. Karataš in M.

Ex affinitate *Archyrophoro uniflora* Bluf. et Fingh. a quo differt caule graciliore, apice vix incrassato et cum foliis setis valde sparsis obsito, capitulis minoribus, involucri phyllis, ligulis, acheniis etc.

A H. Pelivanoviči Petr. in litt. ad Vel. 1887, Fl. bulg. 361 differt caule vix incrassato, valde parce et sparsim setaceo, foliis radicalibus longe oblongo lanceolatis, capitulis mediocribus involucri canescentis phyllis margine ciliatis, ligulis supra tubum dense-, ad dorsum valde parce hirsutis, acheniis scabridis in rostrum breviorē attenuatis.

**Picris stricta** Jord. Cat. Dijon, 1848, p. 19. Hadžibarica pl. et Lubunova in M.

**Tragopogon majus** Jacq. Fl. austr. I, 19, t. 29. M. Korthiati, in coll. ad Oslop et Eksisu in M.

**T. balcanicum** Vel. Fl. bulg. p. 355. In reg. superiore Gorničova pl. M., specimina deflorata cum descriptione l. c. optime congruunt.

**Scorzonia purpurea** L. Sp. pl. 1113. Subsp. *Peristerica* Form. 1898. Radice cylindrica, verticali vel obliqua, collo dense fibroso, caule pumilo 3—8 cm. alto, erecto, semper monocephalo, tenuiter striato, basi pilis crispulis hirsuto, foliis radicalibus lineari-subulatis, 3 nerviis scapo aequilongis vel subbrevioribus, basi interdum pilis crispulis