from Alabama, has oblong leaves with long teeth

and only 2 flowers yellow.

957. CRENAMON Raf. diff. from Sitilias by outer perianthe unequal, pappus plumose, seeds curved compressed tesselated by longitudinal and transverse stria, scapose, corymbose—nearer to Picris than any other G. the name was that of Adanson for Picris, but different by perianthe realy double as in last and seeds not rugose transversaly alone but tesselate, besides habit scapose.

958. Crenamon Pectinatum Raf. smooth, leaves pectinate pinnatifid obtuse, very long and narrow, base attenuate entire, pinnules remote narrow obtuse, scape grooved corymbose 3-6flore, peduncles very long with subulate scales.—Apalachian Mts. Unaka, leaves 6 to 8 inches long, hardly half inch wide, pectinate like the horn of the Saw-Fish, scape pedal, flowers yellow, perianthe acute, outer rather unequal.

959. Krigia petiolaris R. stem flexuose foliose striate pauciflore, leaves all petiolate oblong or obovate minute dentate laciniate, peduncles flliform, perianthes linear—in Alabama, 3 to 6 inches high, leaves only half inch or less, petiol uncial or longer, flowers small few, not

rigidly erect as in Kr. virginica.

960. Krigia integrifolia R. stem erect foliose grooved uniflore, leaves glaucous entire all dissimilar, radical petiolate elliptical obtuse or obovate acute, on stem amplexicaule acute, ovate or lanceolate, perianthe lanceolate—Mts. Alleghanies, how different from last, nearer to Kr. amplexicaulis but smaller with entire leaves, one small flower pale yellow, not orange color. Semipedal, leaves 1 to 3 inches long, all