

Hordeum Hystrix Roth.	Kochia Sedoides Schrad.	Linaria versicolor Moench.	Matricaria pusilla Willd.
— jubatum Linn.	Krigia chilensis Bert. *)	Linum aethiopicum Thunb.	Matthiola fenestralis Cand.
Hornemannia bicolor Willd.	Kruberia leptophylla Hoffm.	— angustifolium Willd.	— parviflora R. Br.
Hyacinthus romanus Linn.	Kuhnia rosmarinifolia Vent.	— austriacum Linn.	— sinuata R. Br.
Hydrocotyle Sibthorpioides Lamk.	Kyllinga monocephala Rottb.	Lippia asperifolia Rich.	— tricuspida R. Br.
Hymenophyllum pubescens C. A. M.		— filiformis Schrad. ***)	Maurandia Barclayana Lindl.
Hyoscyamus orientalis M. B.		— nodiflora Rich.	Medicago brachycarpa Fisch.
— pallidus W. et K.		Lithospermum tenuiflor. Linn. f.	— cancellata M. B.
— pusillus Linn.		Loasa bryoniaefolia Schrad.	— ciliaris Willd.
Hyoseris scabra Linn.	Lactuca ambigua Schrad. **)	— triloba Juss.	— elegans Jacq.
Hypocotyle procumbens Linn.	Lagoecia cuminoidea Linn.	Lobelia heteromalla Schrad.	— maculata Willd.
Hypericum Ascyron Linn.	Lagunaea lobata Willd.	— longiflora Linn.	— Nummularia Cand.
Hyssopus angustifolius M. B.	Lagurus ovatus Linn.	Lolium speciosum M. B.	— praecox Moris.
	Lantana recta Ait.	Lophanthus chinensis Benth.	— uncinata Willd.
	— trifoliata Linn.	— multifidus Benth.	Melanosellum decipiens Hoffm.
	Lasiosperm. pedunculare Lag.	Lophospermum erubescens Don.	Melica Bauhini All.
	— radiatum Trev.	Lotus arabicus Linn.	Melilotus gracilis Trev.
Iberis Lagascana Cand.	Lathyrus articulatus Linn.	— ornithopodioid. Linn.	— longifolia Ten.
— pubescens Willd.	— bithynicus Lamk.	— peregrinus Linn.	— suaveolens Ledeb.
— Tenoriana Cand.	— ciliatus Guss.	— pilosissimus Poir.	Melochia pyramidata Linn.
Impatiens parviflora Cand.	— Ochrus Cand.	Lupinus micranthus Dougl.	Mentha crispata Schrad.
Indigofera caerulea Roxb.	— sphaericus Retz.	— mutabilis Sweet.	— pubescens Willd.
Inula glabra Bess.	— spurius Willd.	— polyphyllus Dougl.	Microtea debilis Sw.
Ipomoea Pes-tigridis Linn.	— tumidus Lamk.	Luzula Forsteri Cand.	Milium multiflorum Cav.
Ixalis aleppica Scop.	Lavatera neapolitana Ten.	— pediformis Cand.	Milleria quinqueflora Linn.
— costata Ledeb.	— plebeja Sims.	— spadicea Cand.	Mimulus alatus Ait.
Ischaemum rugosum Salisb.	Lawsonia inermis Linn.	Lychnis alpina Linn.	— luteus L.
Isolepis curvifolia Schrad.	Lechea minor Linn.	— laeta Ait.	— moschatus Dougl.
— dichotoma Kunth.	Leonitis nepetaefolia R. Br.	— pyrenaica Cand.	— propinquus Lindl.
— Eckloniana Schrad.	Leontodon glaucanthus Ledeb.	Lycopsis pulla Linn.	— rivularis Lindl.
— prolifera R. Br.	Leonurus altaicus Spr.	— rosea Lehm.	— silvestris Hook.
— pubigera Schrad.	— glaucescens Bung.	— versicolor Schrad.	— variegatus H. Par.
— Saviana Schult.	Lepidium affine Ledeb.	— vesicaria Linn.	Mitella diphylla Linn.
Isopyrum sumarioides Linn.	— cordatum Willd.	Lythrum Graefferi Tenor.	Mogiphanes Jacquini Schrad. ****)
Juncus fusco-ater Schreb.	— densiflorum Schrad.		— virgata Schrad.
— monanthos Jacq.	— Eckloni Schrad.		Mollugo verticillata Linn.
— pallidus R. Br.	— Gussoni Schrad.		Monolepis trifida Schrad.
Jussiaea linifolia Vahl.	(Thlaspi pubescens Guss.)		Moscara pinoatifida R. et P.
— ovalifolia Sims.	— incisum Roth.	Madia elegans Don.	Muscari pallens M. B.
Justicia furcata Jacq.	— β. latifol. Ledeb.	— viscosa Cav.	Myosotis versicolor Roth.
— paniculata Burm.	— perfoliatum Linn.	Malachra capitata Linn.	Myrica serrata Willd.
Ixia ramiflora Ten.	— spinosum Linn.	— mexicana Schrad.	
	— suffruticosum Linn.	Malcolmia littorea R. Br.	
	Leucas aspera Link.	— parviflora Cand.	
	— Zeylanica Ait.	Malva angustifolia Cav.	
	Leyssera gnaphaloides Linn.	— borealis Wallm.	Nasturtium chilense H. Par.
Kennedia rubicunda Vent.	Ligusticum scoticum Linn.	— limensis Linn.	— clandestinum Spr.
Kleinia Porrophyllum Linn.	Linaria alpina Willd.	— peruviana Linn.	— erectum Trev.
— suffruticosa Willd.	— reflexa Willd.	— tricuspida Ait.	— pyrenaicum Cand.
Kochia hyssopifolia Roth.	— reticulata Desf.	— walteriaefolia Lk.	Nauenburgia trinervata Willd.
— Scoparia Schrad.	— tristis Willd.	Mariscus aggregatus Willd.	Navarretia pusilla H. Par.

\*) Species anomala ob pappi structuram. Semina ex Hort. Parisino missa.

\*\*) Est media virosam inter et Scariolam, illi praecipue foliis, hinc duratione et habitu gracilliori conveniens. Herba quoque nullum spirat odorem narcoticum, virosae proprium. — Descriptionem et iconem hujus Lactuae dedit Hayne in Plant. Offic. 1. 7. 47. sub virosa.

\*\*\*) f. caule herbaceo repente, foliis oblongo-obovatis basi cuneatis apicem versus serrulatis strigoso-pubescentibus, capitulis ovato-conicis folia longe superantibus, bracteis cuneatis. Chili. 24. Sub diu caules ad bipedalem longitudinem excurrunt, geniculis omnibus radicanibus; nec inferioribus tantum ut in affini nodiflora; folia etiam longius petiolata ac in illa.

\*\*\*\*) Mogiph. Jacquini et virgata, sub brasiliensi Mart. comprehensae, species omnino diversas constituunt sic definiendas: — M. virgata, caule virgato foliisque ovali-oblongis villosis-hirtis, capitulis terminalibus pedunculatis ternis solitariisque, bracteis calice brevioribus. — M. Jacquini, caule ramis patulis foliisque ovatis acuminatis appresso-pubescenti-hirtis, capitulis lateralibus terminalibusque, bracteis interioribus calice longioribus. Ad hanc posteriorem pertinet Gomphrena brasiliensis Jacq., quoad semina ab Ill. Jacquino fil. liberaliter mecum communicata. Icon Jacquini plantam vegetiorem primi anni exhibens, cujus folia duplo majora et paulo longiora his sequentium annorum esse solent.