

rensibus.—*Wight. cat. n.* 1496.—*Carduus indicus*, *Klein!* in herb. *Wight.*—*C. nitidus*, *Wall.!* *cat. n.* 2908.—*Serratula indica*, *Willd. sp. 3. p.* 1642.

### TRIB. VI.—MUTISIACEÆ. *Less. syn. p.* 92.

#### I. DICOMA. *Cass. dict.* 13. *p.* 194; 47. *p.* 503; *Less. syn.* 109.

1. *D. lanuginosa* (DC. prod. :) involuci ovato-oblongi squamis extus glabriusculis, pappi paleis serratis fructu villosissimo vix duplo longioribus.—In pascuis et apricis littoris Coromandeliani.—*Wight. cat. n.* 1497.—*Xeropappus lanuginosus*, *Wall.!* *cat. n.* 2980.—*Acelepis lanata*, *Spreng. in litt. ad Wall.*—*Onopordon lanatum*, *herb. Madr.* (ex *Wall.*)—Valde accedit ad *D. tomentosam*, *Cass.*; sed caract. indicatis facile differt.

### TRIB. VII.—CICHORACEÆ. *Juss. gen.* 168; *Less. syn.* 126.

*Lactuceæ, Cass.*

#### I. CICHIORIUM. *Linn. gen. n.* 921.

1. *C. Cosnia* (Ham. ! in *Wall.!* *cat. n.* 3241:) caule herbaceo erecto laevi hinc inde piloso, foliis oblongis sinuatis aut subpinnatifidis scabridis floralibus basi auriculatis, capitulis axillaribus pluribus congestis sessilibus.—In variis Indiæ locis a Wallichio indicata, sed forte vix indigena ex Wight herb.—*Wight. cat. n.* 1498.—*C. endiviae* affine.

#### II. PICRIS. *Linn. gen. n.* 907.

##### 1. *P. hieracioides* (Linn. sp. *p.* 1114).

*β, indica*; corymbo magis divaricato.—Ad Courtallum.—*Wight. cat. n.* 1499.—Forte *P. hamulosa*, *Wall.!* *cat. n.* 3255.—In Nepaulia lecta ab hac non differt?

#### III. LACTUCA. *Linn. gen. n.* 909; *Less. syn.* 135.

\* *Achænii collo elongato.*

##### 1. *L. sativa* (Linn. sp. 1118).—In Ind. or. culta.—*Wall. cat. n.* 3244.

This is also *L. sativa*, Roxb. fl. Ind. 3. *p.* 403.

\*\* *Achænii collo brevi.* *Mycelis, Cass. dict.*

2. *L. glabra* (DC. prod. :) glaberrima, caule nudo parce ramoso, foliis paulo longiore, foliis elongato-linearibus integrerrimis acutis, capitulis longe pedicellatis cylindraceis subcorymbosis.—In Ind. or.—*Wight. cat. n.* 1500.

3. *L. remotiflora* (DC. prod. :) caule herbaceo tereti glabro subnudo parce ramoso, foliis subradicalibus membranaceis amplexicaulibus, obovatis vix subsinuatis margine setoso-ciliatis cæterum glabris, capitulis secus ramos sub-solitariis distantibus pedicellatis cylindraceis.—In Peninsula indica frequens.—*Wight. cat. n.* 1501.

From a specimen similarly named in Dr Wight's herbarium, this appears to be *Cacalia sonchifolia*, *Wall. L. n.* 3144, d.

4. *L. runcinata* (DC. prod. :) caule erecto glabro tereti apice nudo, foliis subradicalibus rigidulis runcinatis grosse setoso-ciliatis cæterum glabris amplexicaulibus, capitulis secus ramos distantibus subgeminis breviter pedicellatis cylindraceis.—Frequens in ruderatis circa Negapatam.—*Wight. cat. n.* 1502.—*Prenanthes sonchifolia*, *Heyn. in Wall.!* *cat. n.* 5266; *Willd. sp. 3. p.* 1541.—Non est *Lactuca sonchifolia* auct.

*P. racemosa*, Roxb. fl. Ind. 3. *p.* 405, or *Sonchus calyculatus*, Roxb. in E. I. C. mus. tab. 418, belongs to this species.