

linear-oblonga multinervia, obtusa, dentibus marginalibus parvis callosis. *Fl. disci* 30-40, glaberrimi, pappo albo. *Achenia* brevia, costata, glaberrima. *Receptaculus* planiusculum.

A fine species, very nearly allied to the Abyssinian *S. Stuedelii*, Hochst.; but it has more minutely toothed and coriaceous leaves: also allied somewhat to *S. alpina*, *auratus*, &c., but more branched and robust, with sessile, much larger, and differently shaped leaves.

27. *LACTUCA* (*SCARIOLA*) *CAPENSIS*, Thunb., var. *integrifolia*, DC.
Prodr. vii. 136.

Hab. Cameroons Mountains, alt. 5000-7500 feet. (Fl. Nov., Dec.)

Certainly identical with the South-African plant, which extends throughout the Cape Colony to Natal, ascending in the latter country to 5000 feet.

28. *LACTUCA* (*SCARIOLA*) *GLANDULIFERA*, H. f., n. sp. Glandulosa, hispidula, caulibus angulatis flexuosis, ramis apice divaricatis paniculato-ramosissimis, foliis ultra medium ovato-hastatis acutis denticulatis infra medium in petiolum late alatum basi cordato-bilobum abrupte contractis, pedunculis gracilibus chactatis, capitulis 3-4-floris, achenio rostro suo paulo longiore.

Hab. Cameroons Mountains, alt. 7000-8000 feet. (Fl. Dec.-Feb.)

Herba gracilis, scandens, 8-petalis, diffuse ramosa, ramis paniculisque setis glandulosis sparsis. *Folia* 1 unc. longa, membranacea, utrinque pilosula. *Capitula* $\frac{1}{2}$ unc. longa, angusta, squamis extimis brevibus, interioribus e basi ovato-lanceolata longe linearibus apice incrassatis. *Pappus* albus. *Achenia* oblonga, acuta, utrinque medio longitudinaliter anguste 3-5-alata, rostro gracili.

Very near to the Abyssinian *L. paradoxa*, Schultz, in every respect but the glandular setae of the stem and inflorescence.

29. *SONCHUS* *ANGUSTISSIMUS*, H. f., n. sp. Caulis erecto robusto glauco foliisque glaberrimis, foliis e basi lanceolato-hastata anguste elongatis sensim attenuato-acuminatis margine revolutis retrorsum aculeolatis integris v. lobulis 1-2 linearibus recurvis auctis, corymbo umbellato, pedunculis nullis, capitulis setosis basi lanatis.

Hab. Cameroons Mountains, alt. 7000-8000 feet. (Fl. Nov.-Jan.)

Herba 4-6-petalis. *Caulis* erectus, crassit. digiti, striatus, glaucus, simplex. *Folia* spithamea ad pedalis, vix $\frac{1}{2}$ unc. lata, subcoriacea, enervia, costa subtus prominula, lobis basi elongato-subulatis deflexis. *Corymbi* 8-12-cephali, pedunculis 1-3-cephali, basi bracteis setaceis, glabris v. sublanatis, 1 unc. diametro. *Involucri* squamæ lineares, obtusae, basi lana immersae; exteriores setis subflexuosis hirsutae. *Achenia* parva, oblonga, utrinque obtusa, levia, brunnea, compressa, utrinque sub-6-costata, costis obscure transverse undulatis. *Pappus* albus.