

*Achaenium nigrum* valde complanatum oblongum rostris exceptis 4–4½ mm. longum 2½ mm. latum basi obtusum apice acutum ad rostram 1½ mm. longam abeuns facie transverse rugulosum; pappi setæ copiosæ albæ 8 mm. longæ scabræ.

HAB. Mansu (Kōshūn), leg. T. SŌMA, Dec. 1913.

Near *Lactuca brevirostris*, but differs from it in having more elongate heads with narrower bracts.

4. ***Lactuca Morii*** HAYATA sp. nov. (Fig. 30, 31–7). Herba erecta. Caulis 1–2 m. longus toto foliatus subglaber vel tenuiter hirsutus. Folia sessilia pin-




Fig. 30, *Lactuca Morii* HAYATA.

natiloba in ambitu oblonga, lobis lateralibus utroque latere 2–3, irregulariter dentata vel lacerata, lobo terminali minuto lineari caudiv-  
vel cuspidiformi 1½ cm. longo 1 mm. lato, supra scabra subtus pallidissima ad venas costasque setuloso-hirsuta. Capitula racemoso-cymosa laxè disposita. Involucrum late campanulatum 1½ cm. longum totiusque latum basi bracteolis minutis instructum, bracteis involucri exterioribus brevioribus ovatis 1 mm. longis, interioribus longioribus, intimis longissimis linearibus 1½ cm. longis 2 mm. latis glâbris apice obtusis. Fl. ut videntur ochracei. Achaenium maturum nigricans valde complanatum obovatum rostris exceptis 4 mm. longum 2½ mm. latum basi obtusissimum vel rotundatum apice longe rostratum, rostris 3

mm. longis, facie minute transverseque rugulosum; pappi setæ albæ scabræ 6 mm. longæ.

HAB. Hakkudaisan, leg. U. MORI.

Near *L. brevirostris*; but differs from it in having long-cuspidate leaves.

5. ***Lactuca formosana*** MAXIM.; HAYATA l. c. p. 40.

6. ***Lactuca sororia*** MIQ. (Fig. 31–5); HAYATA l. c. p. 40.

HAB. Urai, Agioku, ad 1200 ped. alt., B. HAYATA, Mai. 1916.




Fig. 31; 1, *Lactuca lacerrima* HAY.; 2, *L. taiioensis* HAY.; 3, *L. flavissima* HAY.; 4, *L. longirostra* HAY.; 5, *L. sororia* MIQ.; 6, *L. mansuensis* HAY.; 7, *L. Morii* HAY.; 8, *L. brachyrhyncha* HAY.