

rensis.—*Wight. cat. n. 1496.*—*Carduus indicus, Klein! in herb. Wight.*—*C. nitidus, Wall! cat. n. 2908.*—*Serratula indica, Willd. sp. 3. p. 1642.*

TRIB. VI.—MUTISIACEÆ. *Less. syn. p. 92.*

I. DICOMA. *Cass. dict. 13. p. 194; 47. p. 503; Less. syn. 109.*

1. *D. lanuginosa* (DC. prod. :) involucri ovato-oblongi squamis extus glabrusculis, pappi paleis serratis fructu villosissimo vix duplo longioribus.—In pascuis et apricis littoris Coromandeliani.—*Wight. cat. n. 1497.*—*Xeropappus lanuginosus, Wall! cat. n. 2980.*—*Acelepis lanata, Spreng. in litt. ad Wall.*—*Onopordon lanatum, herb. Madr. (ex Wall.)*—Valde accedit ad *D. tomentosam, Cass.*; sed caract. indicatis facile differt.

TRIB. VII.—CICHORACEÆ. *Juss. gen. 168; Less. syn. 126.*—
Lactucæ, *Cass.*

I. CICHORIUM. *Linn. gen. n. 921.*

1. *C. Cosnia* (Ham! in Wall! cat. n. 3241 :) caule herbaceo erecto lævi hinc inde piloso, foliis oblongis sinuatis aut subpinnatifidis scabridis floralibus basi auriculatis, capitulis axillaribus pluribus congestis sessilibus.—In variis Indiæ locis a Wallichio indicata, sed forte vix indigena ex *Wight herb.*—*Wight. cat. n. 1498.*—*C. endiviæ affine.*

II. PICRIS. *Linn. gen. n. 907.*

1. *P. hieracioides* (Linn. sp. p. 1114).

β, *indica*; corymbo magis divaricato.—Ad Courtallum.—*Wight. cat. n. 1499.*—Forte *P. hamulosa, Wall! cat. n. 3255.*—In Nepaulia lecta ab hac non differt?

III. LACTUCA. *Linn. gen. n. 909; Less. syn. 135.*

* *Achenii collo elongato.*

1. *L. sativa* (Linn. sp. 1118).—In Ind. or. culta.—*Wall. cat. n. 3244.*

This is also *L. sativa, Roxb. fl. Ind. 3. p. 403.*

** *Achenii collo brevi. Mycelis, Cass. dict.*

2. *L. glabra* (DC. prod. :) glaberrima, caule nudo parce ramoso, foliis paulo longiore, foliis elongato-linearibus integerrimis acutis, capitulis longe pedicellatis cylindraceis subcorymbosis.—In Ind. or.—*Wight. cat. n. 1500.*

3. *L. remotiflora* (DC. prod. :) caule herbaceo tereti glabro subnudo parce ramoso, foliis subradicalibus membranaceis amplexicaulibus, obovatis vix subsinuatis margine setoso-ciliatis cæterum glabris, capitulis secus ramos solitariis distantibus pedicellatis cylindraceis.—In Peninsula indica frequens.—*Wight. cat. n. 1501.*

From a specimen similarly named in Dr Wight's herbarium, this appears to be *Cacalia sonchifolia, Wall. L. n. 3144, d.*

4. *L. runcinata* (DC. prod. :) caule erecto glabro tereti apice nudo, foliis subradicalibus rigidulis runcinatis grosse setoso-ciliatis cæterum glabris amplexicaulibus, capitulis secus ramos distantibus subgeminis breviter pedicellatis cylindraceis.—Frequens in ruderalis circa Negapatam.—*Wight. cat. n. 1502.*—*Prenanthes sonchifolia, Heyn. in Wall! cat. n. 5266; Willd. sp. 3. p. 1541.*—Non est *Lactuca sonchifolia* auct.

P. racemosa, Roxb. fl. Ind. 3. p. 405, or Sonchus calyculatus, Roxb. in E. I. C. mus. tab. 418, belongs to this species.

5. *L. napifolia* (DC. prod. :) glabra caulibus erectis nudis ramosis, foliis radicalibus petiolatis lyrato-pinnatilobatis mucronatis lobis oblongis terminali majore, capitulis longe pediculatis parvis, invol. subuniseriali.—In ruderatis circa Negapatam.—*Wight. cat. n. 1503.*—*Prenanthes napifolia, Wall.! herb.*

6. *L. sarmentosa* (Wight herb. !:) glabra, caule flagellifero procumbente hinc inde radices foliaque agente, foliis confertis sinuato-pinnatifidis lobis rotundatis, pedunculis (scapis) 1-cephalis apice bracteolas squamosas subimbricatas margine scariosas gerentibus, invol. cylindraceis.—Frequens in arenosis ad littora maris versus Negapatam.—*Wight. cat. n. 1504.* *Prenanthes sarmentosa, Willd. sp. 3. p. 1540; Wall.! cat. n. 3262.*

The Coromandel specimens mentioned by Roxburgh under *Prenanthes asplenifolia* (fl. Ind. 3. p. 404, or *Hieracium stoloniferum*, Roxb. in E. I. C. mus. tab. 1281), belong to this species.

IV. SONCHUS. *Linn. gen. n. 908.*

1. *S. oleraceus* (Linn. sp. 1115).—Frequens versus Negapatam.—*Wight. cat. n. 1505; Wall.! cat. n. 3252.*

This is *S. oleraceus*, Roxb. fl. Ind. 3. p. 402, or *S. Indicus*, Roxb. in E. I. C. mus. tab. 47; and, judging from Roxburgh's descriptions and drawing, his *S. orivensis*, fl. Ind. 3. p. 402 (in E. I. C. mus. tab. 416), seems also to belong to this variable species.