

latrioribus planis, valvis brevioribus subulatis, foveis circa 15 rotundatis usque 1 mm in diametro,

N. Schoberi a nostra putamine sensim minore (usque 5—7 mm long, 3 mm lat.) foveis profundioribus, valvis longioribus angustis, foliis obtusis, cyma ramis brevibus differt.

N. retusa putamine trigono, acuto, cyma abbreviata, foliis apice regulariter tridentatis vel rotundatis valde differt.

1. V. I. Roborowski, Turkestania Chinensis, in oaso Czerczen 4000'. — 2. J. E. T. Aitchison, no. 724, 27. VII, 1885, Afghanistan, fruct. Haryrud valley. „Comon near Khusan“. Adhuc in Afghanistan, in Turkestania chinensi ad limites meridionales ejus observata est.

CXX. Decades plantarum novarum. XL—XLII.

Auctore H. Léveillé.

(Originaldiagnosen.)

391. *Lactuca nummularifolia* Lévl. et Vant., nov. spec.

Planta acaulis vel breviter caulescens, rosulifera, stolones elongatissimos, graciles et numerosissimos emittens; folia *Nummulariae*, sat longe tamen petiolata, margine remotissime et obscure denticulata; capitula bina, bracteis linearibus; akoenia compressa, apice constricta dein in discum dilatata, pappus candidus.

Corea: Quelpaert in agris Hongno, 15 janv. 1910 (Taquet, 3521).

392. *Salvia Esquirolii* Lévl., nov. spec.

Caulis simplex nudus vel bifoliatus stirpe crasso oriundus, villosus; folia radicalia subcordata rosulata, numerosa, late lanceolata asymmetrica crenata obtusa vel obtusata subtus flavo-ferruginea, supra atroviridia villosa; omnia duobus lobulis liberis basi munita; calyx lutescens, corolla 4-plo brevior, pubescens; corolla rosea vel alba violaceo variegata, stamina labio superiore inclusa; stylus basi villosus, elongatissimus, contortus et bifidus.

Kouy-Tchéou: Se-Tse-Fen, sept. 1904; Lieng-Tchang-Po, juin 1905 (Esquirol, 230, 572).

393. *Salvia tuberifera* Lévl., nov. spec.

S. japonicae var. *bipennatae* affinis; a quo tamen radice conspicue tuberifera, foliis supremis 3-jugis, valerianiformibus, radicalibus autem 3—5 foliolatis, foliolis cordatis rhomboidalibus, distinguitur; corolla parva tubo non exerto calice hirtello dentibus acuminatis vix 2-plo longior; caulis apice tantum et pedicelli glandulosi; flores roseo-violacei.

Kouy-Tchéou: Pien-Yang, avril 1906 (Jul. Cavalerie, 2743).

394. *Salvia betonicoides* Lévl., nov. spec.

Caulis simplex glabrescens; folia radicalia pseudorosulata, caulinaria 2, cordato-oblonga, obtusa (caulinaribus acuminatis) crenata glabrescentia subtus ferruginea; inflorescentia laxe spicata, puberula eglandulosa, nuda;