

ovata, apice acuta, basi cuneata, 1·5–3·3 cm. longa, 1–2 cm. lata, subcoriacea vel rigide chartacea, glabra nisi margine ciliolata, distanter pauci-serrata, trinervata, nervis supra conspicuis subtus prominulis; petioli breves. *Capitula* pluriflora, 2·3 cm. longa; involucri bracteae circiter 6-seriatae, exteriores parvae ovatae, interiores linear-i-oblongae, obtusae, rigidiae, glabrae. *Corollae* tubus 1 cm. longus, superne ampliatus, lobi anguste linear-i-lanceolati, apice revoluti, 0·5 cm. longi. *Antherae* circiter 0·5 cm. longae, basi caudis 2 mm. longis ciliatis instructae. *Achaenia* vix matura, pilosula, costata.

Doi Chieng Dao, 2100 m., *Hosseus*, 405 (Herb. Kew! et Copenhagen!).

*Gerbera piloselloides*, *Cass.*—F.B.I., iii. p. 389; Coll. et Hemsl., Journ. Linn. Soc., xxviii. p. 77.

Chiengmai, on tops of ridges in open jungle on Doi Sootep, 1350–1500 m., *Kerr*, 621.

*Distr.* N. India, China, Trop. and S. Africa, Maserene Islands.

*Crepis acaulis*, *Hook. f.*, F.B.I., iii. p. 396; Coll. et Hemsl., Journ. Linn. Soc., xxviii. p. 78.

Chiengmai, 300 m., *Hosseus*, 442; Doi Sootep, 720 m., *Kerr*, 1727.

*Distr.* India, Burma, Assam, Yunnan.

*Crepis chloroclada*, Coll. et Hemsl., Journ. Linn. Soc., xxviii. p. 78.

Chiengmai, in open grassy jungle on Doi Sootep, 1200–1500 m., *Kerr*, 1108.

*Distr.* Upper Burma, Yunnan.

*Crepis japonica*, *Benth.*—F.B.I., iii. p. 395; Coll. et Hemsl., Journ. Linn. Soc., xxviii. p. 78.

Muang Fang, 300 m., *Hosseus*, 615.

*Distr.* Japan south to Australia and west to Afghanistan and S. Africa.

*Lactuca Parishii*, *Craib*, sp. n., ab affini *L. alatipidi*, Coll. et Hemsl., foliorum lobis numerosioribus, lobo terminali basi fere recto marginibus rectis vel nonnihil concavis sed nunquam convexis, lobis lateralibus a lobo terminali haud tam distantibus, acheniorum rostris multo brevioribus recedit.

*Herba* glabra, robusta, 12 dm. vel ultra alta. *Folia* caulinata inferiore petiolo 7–12 cm. longo alato suffulta, pinnatisecta, 14–31 cm. longa, lobo terminali deltoideo acuto 9–14 cm. longo 7–11 cm. lato, lobis lateralibus utrinque 2–4 supremis a lobo terminali nunquam ultra 1 cm. distantibus plerumque lanceolatis acutis 2–5 cm. longis 1·2–2·5 cm. latis, rachi inter lobos alata, membranacea, margine irregulariter lobata et praeterea distanter denticulata. *Capitula* laxe paniculata, erecta, circiter 12-flora, vix 1·5 cm. longitudinis attingentia. *Achaenia* compressa, 0·5 cm. alta, apice in rostrum 1 mm. longum attenuata, faciebus circa 5-costata, costis superne praecipue pilis brevibus rigidis seabridula.

Burma, Moulmein, 1200–1500 m., *Parish*, 423.

Chiengmai, Doi Sootep, 600–1000 m., *Hosseus*, 454 (Herb. Kew! et Hoss.!), *Kerr*, 1683.