

rensis.—*Wight. cat. n. 1496.*—*Carduus indicus*, *Klein! in herb. Wight.*—*C. nitidus*, *Wall! cat. n. 2908.*—*Serratula indica*, *Willd. sp. 3. p. 1642.*

### TRIB. VI.—MUTISIACEÆ. *Less. syn. p. 92.*

I. DICOMA. *Cass. dict. 13. p. 194; 47. p. 503; Less. syn. 109.*

1. *D. lanuginosa* (DC. prod. :) involucri ovato-oblongi squamis extus glabriusculis, pappi paleis serratis fructu villosissimo vix duplo longioribus.—In pascuis et apricis littoris Coromandeliani.—*Wight. cat. n. 1497.*—*Xeropappus lanuginosus*, *Wall! cat. n. 2980.*—*Acelepis lanata*, *Spreng. in litt. ad Wall.*—*Onopordon lanatum*, *herb. Madr. (ex Wall.)*—Valde accedit ad *D. tomentosam*, *Cass.*; sed caract. indicatis facile differt.

### TRIB. VII.—CICHORACEÆ. *Juss. gen. 168; Less. syn. 126.*— Lactuceæ, *Cass.*

I. CICHORIUM. *Linn. gen. n. 921.*

1. *C. Cosnia* (Ham! in Wall! cat. n. 3241 :) caule herbaceo erecto lævi hinc inde piloso, foliis oblongis sinuatis aut subpinnatifidis scabridis floralibus basi auriculatis, capitulis axillaribus pluribus congestis sessilibus.—In variis Indiæ locis a Wallichio indicata, sed forte vix indigena ex *Wight herb.*—*Wight. cat. n. 1498.*—*C. endiviæ* affine.

II. PICRIS. *Linn. gen. n. 907.*

1. *P. hieracioides* (Linn. sp. p. 1114).

*β, indica*; corymbo magis divaricato.—Ad Courtallum.—*Wight. cat. n. 1499.*—Forte *P. hamulosa*, *Wall! cat. n. 3255.*—In Nepaulia lecta ab hac non differt?

III. LACTUCA. *Linn. gen. n. 909; Less. syn. 135.*

\* *Achenii collo elongato.*

1. *L. sativa* (Linn. sp. 1118).—In Ind. or. culta.—*Wall. cat. n. 3244.*

This is also *L. sativa*, *Roxb. fl. Ind. 3. p. 403.*

\*\* *Achenii collo brevi.* *Mycelis, Cass. dict.*

2. *L. glabra* (DC. prod. :) glaberrima, caule nudo parce ramoso, foliis paulo longiore, foliis elongato-linearibus integerrimis acutis, capitulis longe pedicellatis cylindræis subcorymbosis.—In Ind. or.—*Wight. cat. n. 1500.*

3. *L. remotiflora* (DC. prod. :) caule herbaceo tereti glabro subnudo parce ramoso, foliis subradicalibus membranaceis amplexicaulibus, obovatis vix subsinuatis margine setoso-ciliatis cæterum glabris, capitulis secus ramos subsolitariis distantibus pedicellatis cylindræis.—In Peninsula indica frequens.—*Wight. cat. n. 1501.*

From a specimen similarly named in Dr Wight's herbarium, this appears to be *Cacalia sonchifolia*, *Wall. L. n. 3144, d.*

4. *L. runcinata* (DC. prod. :) caule erecto glabro tereti apice nudo, foliis subradicalibus rigidulis runcinatis grosse setoso-ciliatis cæterum glabris amplexicaulibus, capitulis secus ramos distantibus subgeminis breviter pedicellatis cylindræis.—Frequens in ruderatis circa Negapatam.—*Wight. cat. n. 1502.*—*Prenanthes sonchifolia*, *Heyn. in Wall! cat. n. 5266; Willd. sp. 3. p. 1541.*—Non est *Lactuca sonchifolia* auct.

*P. racemosa*, *Roxb. fl. Ind. 3. p. 405*, or *Sonchus calyculatus*, *Roxb. in E. I. C. mus. tab. 418*, belongs to this species.