

4. LACTUCA. *Linn.* (*part of.*)

*Involucrum cylindraceum*, squamis seu foliolis erectis imbricatis, exterioribus latioribus. *Receptaculum* planum, nudum. *Achenia compressa*. *Pappus* niveus, stipitatus, pilis scabris seu laevibus.

1. *L. elongata*; subglabra, foliis uncinatis dentatis subtus glaucis inferioribus petiolisque hirsutis superioribus subintegris, floribus corymboso-paniculatis. *Willd. Enum.* p. 817. *Pursh, Fl. Am.* v. 2. p. 500. *Elliott, Carol.* v. 2. p. 252.—*L. longifolia*. *Mich. Am.* v. 2. p. 85. (fide Elliott.)—*L. hirsuta*. *Muhl. Cat.*—*Nutt. Nov. Gen.* v. 2. p. 124?

HAB. Lake Huron. *Dr. Todd*. Plains of the Saskatchewan. *Dr. Richardson*. *Drummond*.—As the lower leaves of this plant are frequently hairy, especially upon the petioles, I am disposed to believe that the *L. hirsuta* of Muhl. Cat. is the same species, and Dr. Torrey entertains a similar opinion.

5. LEONTODON. *Linn.*

*Involucrum imbricatum*, foliolis appressis seu exterioribus patentibus. *Receptaculum* planum, punctatum, nudum. *Pappus* stipitatus, simplex.

1. *L. Taraxacum*; involuci squamis inferioribus reflexis, foliis runcinatis dentatis.—*Linn. Sp. Pl.* p. 1122. *Engl. Bot. t.* 510. *Pursh, Fl. Am.* v. 2. p. 497.

HAB. Throughout the British Possessions, and on both sides of the Rocky Mountains.

2. *L. palustre*; involuci foliolis erectis appressis, foliis sinuato-dentatis glabriusculis.—*Engl. Bot. t.* 533. *Rich. in Frankl. 1st Journ. ed. 2. App.* p. 29.

HAB. Throughout the British Possessions, and from Hudson's Bay to the Pacific.

3. *L. hirsutum*; ubique hirsutum, involuci foliolis erectis appressis, foliis pinnatifidis segmentis inæqualibus remotis acuminatis.

HAB. Menzies Island, and sandy banks of the Columbia. *Douglas*. *Dr. Scouler*.—The leaves of this are very different from any of the numerous varieties of the two preceding species that I have seen; and they and the scapes and involucres are very hairy; the latter clothed with fulvous hairs.

6. APARGIA. *Schreb.*

*Involucrum imbricatum*, foliolis interioribus æqualibus, exterioribus minoribus. *Receptaculum* nudum, punctatum. *Pappus* plumosus, sessilis.

1. *A. autumnalis*; scapo ramoso superne squamoso, foliis lanceolatis dentatis seu pinnatifidis glabriusculis, pedunculis superne incrassatis, involucro subpubescente.—*Willd. Sp. Pl.* p. 1550. *Pursh, Fl. Am.* v. 2. p. 497.—*Hedypnois autumnalis*. *Huds.*—*Engl. Bot. t.* 830.—*Leontodon autumnale*. *Linn.*

HAB. Newfoundland. *Mr. Cormack*.—Pursh gives this plant as a doubtful native of the New England States. It is probably also introduced into Newfoundland from Europe.

## 7. LAPSONA.

*Involucrum subcylindraceum* e foliolis paucis basi squamu-  
losum. *Receptaculum* nudum. *Achenia oblonga* basi attenuata. *Pappus nullus*.

1. *L. communis*; involuci fructu angulato, caule paniculato, pedunculis gracilibus, foliis ovatis petiolatis angulato-dentatis.—*Linn. Sp. Pl.* p. 1141. *Engl. Bot. t.* 844.

HAB. Canada. *Mrs. Sheppard*. Probably introduced from Europe.