

with large rayed heads and dark brown strongly fringed phyllaries, which seems to be the †*C. pratensis* Thuill. of the *Flore de Normandie*. — *C. paniculata* L. Still in the same locality at St. Ouen's, where I saw it in 1877; this year it was much dwarfed by the drought. — *C. aspera* L. Abundant at St. Ouen's, and I think native; some of the capitula had yellowish-white flowers. Also at Vazon Bay, Guernsey, and in Alderney. — *C. Cyanus* L. On the walls at St. Peter's Port, Guernsey; St. Aubin's, Jersey.

*Cichorium Intybus* L. St. Martin's, Jersey.

*Hypochæris glabra* L. Bel Royal, Don Bridge, &c., Jersey. —

*H. radicata* L. Still near Grosnez; it was in beautiful flower. —

†Var. *hispida* Peterm. St. Helier's; Alderney.

*Picris hieracioides* L. †var. *incana* mihi. Between the town and Longy Bay by the roadside in considerable quantity. This occurred as a well-marked variety with narrow leaves deeply and unevenly cut, with densely aggregated capitula, the centre ones being much overtopped by the lateral ones. The outer phyllaries covered with a short greyish-white tomentum, resembling that of *Crepis taraxacifolia*. — †Var. *arvalis* (Jord.). Near Pont Marquet, Jersey.

*Leontodon hispidus* L. Babington says it is common in the Channel Isles, but neither Mr. Lester-Garland nor I was able to find this in the islands. — *L. nudicaulis* (L.) ? Mérat (*L. hirtum* L.). Very common in Jersey and Guernsey, and very variable. — †Var. *arenaria* (DC.) (sub *Thrincia hirta*). A more glabrous and much branching plant, with strong tap-root and leaves sinuately cut, with hairy phyllaries, grows in the sands at St. Helier's and St. Aubin's, Jersey; near Vazon, Guernsey; and at Longy Bay, Alderney. — †Var. *pristis* mihi. This is a densely-hispidly hairy, subsimple plant, with strong tap-root, the leaves deeply repand-runcinate and covered with long shaggy hairs at the base. The phyllaries are subglabrous. I saw this at Vale, Guernsey, and at Longy Bay, Alderney. I think it is the *L. hispidus* of Babington, but the fruit-characters agree with *Thrincia*. The plant looks specifically distinct from *nudicaulis*. The name is given from the leaf-cutting suggestive of the saw-fish. Another plant with long narrow, nearly entire, leaves and glabrous phyllaries, which grew at Vale, Guernsey, is the †var. *leiolæna* (Bisch.).

*Sonchus asper* Hill var. *runcinatus*. Mont Orgeuil, Jersey.

*Crepis capillaris* Wallr. A form of this plant was plentiful by St. Aubin's Bay, and on the Quenvais, Jersey, and also in Alderney, which had a strong tap-root, with numerous flowers of a deep yellow colour, the outer ligules being tipped with red; the plants from four to six inches high.

*Tragopogon porrifolius* L. Near the harbour, Alderney, quite naturalized, and fruiting freely. Casual at St. Luke's, Jersey.

†*Symphytum asperum* Lepechin (*asperrimum* Donn), the fodder plant. I saw it at St. Catherine's and St. Brelade's, Jersey.

*Anchusa sempervirens* L. St. Aubin's, &c. Apparently native in Jersey.

*Myosotis scorpioides* L. (= *M. arvensis* Lam.). Pont Marquet,