

Holotype: Frequent along the shingle beach, Snettisham Scalp, W. Norfolk, v.c. 28, 53/649339, 15 July 1987, P. D. Sell no. 87/213 and L. C. Nicol (**CGE**).

*Caules* plerumque breviores quam 50 cm. *Capitula* 1–3, in ramis brevibus portata.

*Cirsium vulgare* (Savi) Ten., *Fl. Napol.* **5**: 209 (1835–1836).

var. **litorale** P. D. Sell var. nov.

Holotype: Frequent along the shingle beach, Snettisham Scalp, W. Norfolk, v.c. 28, 53/649339, 15 July 1987, P. D. Sell no. 87/211 and L. C. Nicol (**CGE**).

*Caules* ad 50 cm. *Capitula* pauca, in ramis brevibus congesta.

*Centaurea scabiosa* L., *Sp. Pl.* 913 (1753).

var. **nana** P. D. Sell var. nov.

Holotype: Exposed grassy bank above the sea, near Port Eynon, Gower Peninsula, Glamorganshire, v.c. 41, 21/467844, 2 May 1976, P. D. Sell no. 76/48 (**CGE**).

*Planta* ad 20 cm. *Folia* profunde pinnatisecta. *Capitula* pauca, in pedunculis brevibus portata.

*Centaurea debeauxii* Gren. & Godr., *Fl. France* **2**: 243 (1850).

subsp. *nemoralis* (Jord.) Dostál in *Bot. Jour. Linn. Soc.* **71**: 207 (1976).

var. **minima** (C. E. Britton) P. D. Sell comb. nov.

*C. nemoralis* var. *minima* C. E. Britton in *Rep. Bot. Soc. Exch. Cl. Brit. Isles* **6**: 414 (1922).

var. **nemoralis** (Jord.) P. D. Sell comb. nov.

*C. nemoralis* Jord., *Mém. Acad. Sci. Lyon* **1** (1851) (*Pug. Pl. Nov.* 104 (1852)).

*C. nigra* var. *nemoralis* (Jord.) Greml., *Exkursionsfl. Schweiz* ed. 6, 254 (1889).

subsp. *thuillieri* Dostál in *Bot. Jour. Linn. Soc.* **71**: 207 (1976).

var. **pumila** P. D. Sell var. nov.

Holotype: Freshwater Downs, Isle of Wight, August 1890, J. H. A. Stewart (**CGE**).

*Planta* ad 30 cm. *Capitula* pauca, in pedunculis brevibus portata. *Flores exteriore*s interioribus multo longiores.

var. **thuillieri** (Dostál) P. D. Sell stat. nov.

*C. debeauxii* subsp. *thuillieri* Dostál in *Bot. Jour. Linn. Soc.* **71**: 207 (1976).

*C. pratensis* Thuill., *Fl. Env. Paris* ed. 2, 444 (1799), non Salisb., *Prod. Stirp. Chap. Allerton* 208 (1796).

*Scorzoneroïdes autumnalis* (L.) Moench, *Methodus Pl.* 549 (1794).

*Leontodon autumnalis* L., *Sp. Pl.* 798 (1753).

var. **salina** (Aspegren) P. D. Sell comb. nov.

*Apargia autumnalis* var. *salina* Aspegren, *Blekingsk Fl.* 57 (1823).

var. **simplex** (Duby) P. D. Sell comb. nov.

*Leontodon autumnalis* var. *simplex* Duby, *Bot. Gall.* ed. 2, **1**: 308 (1828) pro Sowerby, *Eng. Bot. tab.* 830 (1801).

var. **nigrolanata** (Fr.) P. D. Sell comb. nov.

*Leontodon autumnalis* var. *nigrolanatus* Fr., *Summa Veg. Scand.* 8, 187 (1846).

var. **alpina** (Gaudin) P. D. Sell comb. nov.

*Apargia autumnalis* var. *alpina* Gaudin, *Fl. Helv.* **5**: 59 (1829).

var. **pratensis** (Hornem.) P. D. Sell comb. et stat. nov.

*Apargia pratensis* Hornem., *Hort. Bot. Hafn.* **2**: 785 (1815).

var. **pinnatifida** (Schur) P. D. Sell comb. nov.

*L. autumnalis* var. *pinnatifidus* Schur, *Enum. Pl. Transsilv.* 356 (1866).

var. **coronopifolia** (Lange) P. D. Sell comb. nov.

*L. autumnalis* var. *coronopifolius* Lange, *Haandb. Danske Fl.* ed. 4, 399 (1886).

var. **cinerascens** (Briq.) P. D. Sell comb. nov.

*L. autumnalis* var. *cinerascens* Briq. in *Annu. Cons. Jard. Bot. Genève* **3**: 124 (1899).

var. **dentata** (Holuby ex Schur) P. D. Sell comb. nov.

*L. autumnalis* var. *dentatus* Holuby ex Schur in *Verhandl. Naturf. Ver. Brünn* **15**: 186 (1877).

var. **latifolia** (Schur) P. D. Sell comb. nov.

*L. autumnalis* var. *latifolius* Schur, *Enum. Pl. Transsilv.* 356 (1866).

**Leontodon × vegetus** Finch & P. D. Sell nothospecies nova  
= **L. hispidus × saxatilis**

Holotype: Origin: Parish Church (St James the Less), North Lancing, W. Sussex, v.c. 13, 51/182055, 5 July 1996, P. D. Finch. Growing with both parental species nearby, in turf just outside the kerb at east end of graves of the Dabbs family, to the south of church, west of path down to Vicarage. Cult. in Lancing soil with coco-peat at bottom of pots till pressed on 29 August 1997, by R. A. Finch.  $2n = 11$ .  $F_1$ . (**CGE**).

*Herba perennis* variabilis, praesertim magnitudine, induimento, foliorum figura, papporum auctu. *Caules* ad 45 cm, graciles. *Folia* ad  $20.0 \times 2.5$  cm, pallide luteoviridia, oblongooblanceolata, ad apicem obtusum angustata, sinuatodentata, pilis diverse furcatis vestita, ad basin in petiolum gracilem longe attenuata. *Involucri bracteae* ad  $11 \times 2$  mm, pallide virides, glabrae vel pilis numerosis diverse furcatis pallidis vestitae. *Cypselae* pleraeque steriles maleque effeciae, exteiiores plerumque pappo ad instar coronae, interiores pilorum plumosorum pappo praeditae. *Pollinis grana* magnitudine variabilis.

**Leontodon saxatilis** Lam., *Fl. Franç.* **2**: 115 (1779).

var. **pristis** (Druce) P. D. Sell comb. nov.

*L. nudicaulis* var. *pristis* Druce in *Jour. Bot. (London)* **45**: 423 (1907).

var. **arenarius** (Duby) P. D. Sell comb. nov.

*L. hastilis* var. *arenarius* Duby, *Bot. Gall.* ed. 2, **1**: 302 (1828).

*Picris hieracioides* L., *Sp. Pl.* 792 (1753).

var. **villarsii** (Jord.) P. D. Sell comb. nov.

*P. villarsii* Jord., *Cat. Dijon* 29 (1848).

var. **leteae** (E. I. Nyár.) P. D. Sell comb. nov.

*P. sonchoides* var. *leteae* E. I. Nyár. in *Fl. Rep. Pop. Rom.* **10**: 719 (1965).

var. **arenaria** (E. I. Nyár.) P. D. Sell comb. nov.

*P. sonchoides* var. *arenaria* E. I. Nyár. in *Fl. Rep. Pop. Rom.* **10**: 720 (1965).