

4. { Involucri phylla 7 mm. longa, caulis basi hirsutissimus. — Washington, ad Mason City (*Sandberg!*).  
**Hieracium candelabrum** Gdgr.
4. { Involucri phylla 10 mm. longa, caulis basi glaber. — Wyoming, ad Centennial Valley (*Nelson n. 1678!*).  
**Hieracium Nelsoni** Gdgr.
1. { Folia superiora pinnata vel incisa, rami patuli pauciflori, phylla intima involucri 8 mm. longa. — America bor., Texas (*Lindheimer an. 1849-51!*) ..... **Mulgedium Lindheimeri** Gdgr.
1. { Folia sup. pinnata, rami adscendentes multiflori, phylla intima involucri 11 mm. longa. — Amer. bor., Kansas ad Riley (*Norton n. 315!*)..... **Mulgedium polyanthum** Gdgr.
1. { Folia sup. integerrima, rami uniflori, phylla intima 9,5 mm. longa. — America bor., Illinois ad West Pullman (*Lansing n. 668!*)..... **Mulgedium centrale** Gdgr.

Tres præcedentes, quoad habitum, etc., sat dissimiles, prope *M. floridanum* DC. sunt collocandæ.

✓ **Microderis nevadensis** Gdgr. — Biennis prorsus papilloso-canescens ad collum araneosa, folia nervosa primatifida fidis lanceolatis valde ciliatis, pedunculi puberuli tubo incrassati, achenia apice tantum hirtella paleis profunde bifidis.

HAB. : California, Nevada ad Reno (*Doten! Kennedy! Heizer!*).

A *M. linariifolia* Sch. bip. (*Suksdorf n. 5583!*), etc., differt indumento, foliis magis sectis.

*Microderis Forsteri* Hook. f. in Oceania multum variat; e sequentibus conjice :

- |    |   |                                                                                                                                                                                           |   |
|----|---|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---|
| 1. | { | Folia incisa vel subpinnatifida.....                                                                                                                                                      | 2 |
|    | { | Folia integerrima.....                                                                                                                                                                    | 5 |
| 2. | { | Folia obtusa.....                                                                                                                                                                         | 3 |
|    | { | Folia lanceolata.....                                                                                                                                                                     | 4 |
| 3. | { | Scapus glaber, phylla intima involucri ligulis subbreviora. — Tasmania, Little Plains ( <i>Simson n. 527!</i> ).<br><b>Microderis obtusifolia</b> Gdgr.                                   |   |
|    | { | Scapus pulverulentus, phylla intima involucri ligulis triplo breviora. — Tasmania, ad Sandy Bay ( <i>Spicer n. 94! ex parte</i> ).<br><b>Microderis tenuicula</b> Gdgr.                   |   |
| 4. | { | Calyculus purpureus, folia lata undulata, scapus glaber. — Australia, N. S. Wales ad Tia Falls ( <i>Forsyth!</i> ).<br><b>Microderis latifolia</b> Gdgr.                                  |   |
|    | { | Calyculus viridis, folia angusta plana, scapus fere toto pulverulentus. — Tasmania, ad Kangaroo Point ( <i>Spicer n. 94 p. p.!</i> )<br><b>Microderis tasmanica</b> Gdgr.                 |   |
| 5. | { | Folia lanceolata nervosa 2/3 sup. scapi glabri æquantia, capitula late cylindrico-ovata. — Australia, Victoria ad Sandringham ( <i>Walter!</i> ) ..... <b>Microderis Walteri</b> Gdgr.    |   |
|    | { | Folia obtusa vix nervosa 1/3 inf. scapi pulverulenti æquantia, cap. anguste cylindrico-tubulosa. — Nova Zelandia ad Dunedin ( <i>Thomson!</i> )..... <b>Microderis neozelandica</b> Gdgr. |   |