

4. { Involucri phylla 7 mm. longa, caulis basi hirsutissimus. — Washington, ad Mason City (*Sandberg!*).
Hieracium candelabrum Gdgr.
4. { Involucri phylla 10 mm. longa, caulis basi glaber. — Wyoming, ad Centennial Valley (*Nelson n. 1678!*).
Hieracium Nelsoni Gdgr.
1. { Folia superiora pinnata vel incisa, rami patuli pauciflori, phylla intima involucris 8 mm. longa. — America bor., Texas (*Lindheimer an. 1849-51!*) **Mulgedium Lindheimeri** Gdgr.
1. { Folia sup. pinnata, rami adscendentes multiflori, phylla intima involucris 11 mm. longa. — Amer. bor., Kansas ad Riley (*Norton n. 315!*)..... **Mulgedium polyanthum** Gdgr.
1. { Folia sup. integerrima, rami uniflori, phylla intima 9,5 mm. longa. — America bor., Illinois ad West Pullman (*Lansing n. 668!*)..... **Mulgedium centrale** Gdgr.

Tres præcedentes, quoad habitum, etc., sat dissimiles, prope *M. floridanum* DC. sunt collocandæ.

✓ **Microderis nevadensis** Gdgr. — Biennis prorsus papilloso-canescens ad collum araneosa, folia nervosa primatifida fidis lanceolatis valde ciliatis, pedunculi puberuli tubo incrassati, achenia apice tantum hirtella paleis profunde bifidis.

HAB. : California, Nevada ad Reno (*Doten! Kennedy! Heizer!*).

A *M. linariifolia* Sch. bip. (*Suksdorf n. 5583!*), etc., differt indumento, foliis magis sectis.

Microderis Forsteri Hook. f. in Oceania multum variat; e sequentibus conjice :

- | | | | |
|----|---|---|---|
| 1. | { | Folia incisa vel subpinnatifida..... | 2 |
| | { | Folia integerrima..... | 5 |
| 2. | { | Folia obtusa..... | 3 |
| | { | Folia lanceolata..... | 4 |
| 3. | { | Scapus glaber, phylla intima involucris ligulis subbreviora. — Tasmania, Little Plains (<i>Simson n. 527!</i>).
Microderis obtusifolia Gdgr. | |
| | { | Scapus pulverulentus, phylla intima involucris ligulis triplo breviora. — Tasmania, ad Sandy Bay (<i>Spicer n. 94! ex parte</i>).
Microderis tenuicula Gdgr. | |
| 4. | { | Calyculus purpureus, folia lata undulata, scapus glaber. — Australia, N. S. Wales ad Tia Falls (<i>Forsyth!</i>).
Microderis latifolia Gdgr. | |
| | { | Calyculus viridis, folia angusta plana, scapus fere toto pulverulentus. — Tasmania, ad Kangaroo Point (<i>Spicer n. 94 p. p.!</i>)
Microderis tasmanica Gdgr. | |
| 5. | { | Folia lanceolata nervosa 2/3 sup. scapi glabri æquantia, capitula late cylindrico-ovata. — Australia, Victoria ad Sandringham (<i>Walter!</i>) Microderis Walteri Gdgr. | |
| | { | Folia obtusa vix nervosa 1/3 inf. scapi pulverulenti æquantia, cap. anguste cylindrico-tubulosa. — Nova Zelandia ad Dunedin (<i>Thomson!</i>)..... Microderis neozelandica Gdgr. | |