

Zhejiang (浙江): Xuexi (雪栖), S. Y. Zhang (章绍尧) 972.

### 2. 三裂假福王草 新种

**Paraprenanthes multiformis** Shih, sp. nov.

Affinis speciei praecedenti, quae foliis indivisis et caulis et ramis glabris distinguitur.

Planta annua. Caulis solitarius, 40—90 cm altus, basi 4 mm in diametro, superne paniculato-ramosus, ramis gracilibus, villis multicellularibus noduliformibus obtectis. Folia basalia ignota; folia inferiora mediaque triangulato-vel ovato-hastata vel longe cordata, 6—8 cm longa, 5—6 cm lata, trilobata vel tripatis, lobis omnibus margine denticulatis vel apiculato-dentatis, apice acutis; folia superiora mediis similia, sed minora vel indivisa, triangulata, margine apiculato-denticulata; folia suprema indivisa, lanceolata, triangulato-lanceolata vel anguste elliptica, apice longe acuminata, margine apiculato-denticulata vel subintegrima; omnia longe petiolata, petiolis 3—10 cm longis, interdum late alatis, textura crassiuscula, utrinque aspera, strigoso-furfuracea. Calathidia numerosa, ad caulem et ramos in inflorescentiam paniculatam terminalem disposita. Involucrum cylindricum, calyculatum, 1 cm longum, c. 3 mm in diametro, post anthesin non ovoideum, phyllis 2(3)-seriatis, extimis minimis, lanceolatis, 1—2 mm longis, vix 1 mm latis, apice acutis, intimis longissimis, linearis-lanceolatis vel longe ellipticis, 1 cm longis, 1 mm latis, apice obtusis, omnibus glabris, pallide purpureis. Flosculi ligulati, rosei, c. 11 in numero, ligulis apice minute 5-fidis, tubis ad faucem sparse albo-pilosus. Achaenia atrata, fusiformia, crassiuscula, 4 mm longa, 0.8 mm lata, superne breviter attenuata, apice albida, erostrata, facie 6-costulata. Pappi setae 2-seriatae, 6.5 mm longae, obscure minute scabridae.

Hunan (湖南): Zhixian (芷县), in sylva, alt. 800m, 28 VI 1859, P. X. Tan (谭沛祥) 61587 (Typus, PE). Jiangxi (江西): sine loco, in sylva, Jiangxi Exped. (江西队) 656. Sichuan (四川): Emeishan (峨眉山), ad marginem sylvae, alt. 600 m, Y. L. Chen (陈艺林) 006. Fujian (福建): Shunchang (顺昌), in declivio alt. 500m, Zh. Y. Li (李振宇) 5209.

### 3. 箭耳假福王草 新种

**Paraprenanthes sagittiformis** Shih, sp. nov.

Affinis *P. auriculiformi* Shih, a qua foliis saltem basalibus lyrato-pinnatipartitis, basi sagittiforme semiamplexicaulibus bene differt.

Herba perennis, rhizomate elongato, tenui, prostrato. Caulis erectus, 80 cm altus, superne anguste paniculato-ramosus, ramis gracilibus, glabris. Folia basalia lyrato-pinnatisecta, petiolata, petiolis 5 cm longis, lobis terminalibus triangulatis, 6 cm longis, 5.5 cm latis, lobis lateralis 1-jugis, ellipticis, 1.5 cm longis, 1 cm latis, basi confluentibus; folia inferiora mediaque pinnatipartita, ambitu oblanceolata, lanceolata vel longe elliptica, 10—20 cm longa, 5—9 cm lata, sessilia, basi sagittiforme semiamplexicaulia, lobis terminalibus longe lanceolatis vel ellipticis, 7—9 cm longis, 1—2 cm latis, apice acuminatis, margine minute apiculato-denticulatis, lobis lateralis (1)3—5-jugis, oppositis, oblique oppositis vel alternatis, rhombiformibus, oblique rhombiformibus, ellipticis vel longe ellipticis, apice acutis, margine apiculato-dentatis veldenticulatis, supremis 2.5—4 cm longis, 1—2 cm latis, infimis minimis, auriculiformibus; folia superiora et suprema anguste linearia vel linearis-lanceolata, indivisa, minora, apice longe acuminata, margine minute apiculato-denticulata vel apiculata, basi cuneata, sessilia; omnia