

In *Sonchus* the rays of the pappus are united into bundles at the base, and the receptacle is rough and much dilated.

Trib. 4. SCORZONERÆ. *Receptaculum* epaleatum. *Antheræ* basi bisetæ, appendiculâ exiguâ reniformi! coronatæ. *Stigmata* sæpiùs filiformia, papillosa. *Involucrum* simplex v. imbricatum.

#### PICROSTIA.

*Involucrum* 8-phyllum. *Ligulæ* tridentatæ, 4-nerviæ! *Pappus* capillaris, stipitatus.

*Involucrum* simplici serie 8-phyllum, cylindricum, basi omnino nudum: *foliolis* ligulatis, obtusis, margine invicem se imbricatis, planis. *Receptaculum* planum, nudum. *Flosculi* plurimi: *tubus* tenuissimus, filiformis; *fauce* extùs villosiusculâ: *ligulâ* linearî, tridentatâ, 4-nerviâ. *Stamina* ferè omnino exserta: *filamenta* capillaria, libera: *antheræ* leviter coalitæ, basi bidenticulatae, cristâ exiguâ brevissimâ coronatæ. *Stigmata* brevia, semicylindrica, obtusa, papilloso-pruinosa, arcuato-conniventia. *Achenia* fusiformia, sulcis plurimis longitudinaliter notata, glabra. *Pappus* longè stipitatus, capillaris, mollis, caducus: *radiis* duplice ordine confertis, scabriusculis, inæqualibus, basi solutis.

Herba perennis, lœvigata, glauca. Caulis erectus, pedalis, ramosus, teres, striatus, calatum scriptorium crassitie ferè adæquans. Rami paucissimi. Folia sessilia, lanceolata, integriflora, nunc remotissimè dentata, palmaria v. spithamea, pollicem lata; inferiora basin versus angustata, sublingulata; superiora amplexicaulia, basi sagittata. Flores solitarii, pendunculati. Involucrum unciale, lœvissimum: foliolis apice puberulis. Flosculi lutei? tenues, involucro longiores. Pappus semipollicem longus, cinereo-fulvus. Nomen a πικρος, amarus.