

20. *Lactuca*, sp. n. *L. stoloniferæ* aff.; gracillima, longe stolonifera, foliis petiolatis, cordato-rotundatis, paucilobatis simul calloso-denticulatis.

HUPEH: Patung district (*A. Henry!*). Herb. Kew.

Apparently a very distinct undescribed species; but as the specimens are exceedingly young and otherwise imperfect, it is impossible to characterize it.

21. *Lactuca*, sp. n.? Aff. *L. triangulatæ*, differt achæniis atris utrinque 5-6-costatis.

SHINGKING: without locality (*Ross!*). Herb. Kew.

This has a long narrow inflorescence, and is probably an undescribed species; but the specimen bears no perfect leaves.

1. *Prenanthes Faberii*, *Hemsl.*, n. sp.

*Herba* undique glabra vel cito glabrescens, caulibus crassiusculis. *Folia* caulina petiolata, papyracea, discoloria, deltoideo-cordiformia, usque ad 6 poll. diametro, obscure paucilobata, simul remote calloso-denticulata, lobis acutiusculis vel acuminatis, venis palmato-radiatis subtus sat conspicuis; petiolus gracilis, 1-2 poll. longus. *Capitula* sæpissime 5-flora, interdum usque ad 10-flora, anguste racemoso-paniculata, breviter pedunculata; involucri bracteæ herbaceæ, latæ, obtusæ, extimæ breves, intimæ sæpius 5 pappo breviores; receptaculum nudum. *Florum* color? *Achænia* (immatura tantum visa) brevia, truncata, glabra; pappi setæ numerosissimæ, fulvæ, minute barbellatæ.

SZETCHUEN: Mount Omei, at 10,000 feet (*Faber!*). Herb. Kew.

This differs from *P. Tatarinowii*, its nearest ally, in being glabrous, in the thicker flower-heads, in short dense racemes, forming a long narrow panicle.

2. *Prenanthes ochroleuca*, *Hemsl.*

*Nabalus ochroleucus*, *Maxim. in Bull. Acad. Pétersb.* xv. 1871, p. 376; *Franch. et Savat. Enum. Pl. Jap.* i. p. 274.

COREA: Chemulpo (*Carles!*). Herb. Kew.

Japan.

3. *Prenanthes Tatarinowii*, *Maxim. Ind. Fl. Pek. in Prim. Fl. Amur.* p. 474 in adnot., et in *Mél. Biol.* ix. p. 365; *Franchet, Pl. David.* p. 189.