

598. PRENANTHES. Gen. pl. 1236.

juncea.

1. P. caule virgatim-ramosissimo sulcato glabro, foliis caulinis remotis subulatis brevissimis, ramulis unifloris, calycibus 5-fidis 5-floris: foliolis margine membranaceis.

On the banks of the Missouri. 24. v. s. Flowers purple; it has a striking resemblance to *Chondrilla juncea*; but the leaves are shorter, and the pappus entirely sessile; the radical leaves I have not seen.

altissima.

2. P. caule ramoso, foliis trilobis petiolatis angulatis denticulatis margine scabris, racemis axillaribus, floribus nutantibus, calycibus sub-5-floris.—Willd. sp. pl. 3. p. 1537.

Icon. Pluk. alm. t. 317. f. 2.

In shady woods: Canada to Virginia. 24. July—Sept. v. v. Flowers yellow. Generally a very tall robust plant.

cordata.

3. P. caule superne paniculato, foliis petiolatis cordatis dentatis ciliatis, floralibus sessilibus oblongis integerrimis, panicula laxa racemiflora, floribus nutantibus, calycibus 6-fidis 6-floris.—Willd. hort. berol. 25.

In shady woods, about rocks; New York to Virginia.

24. Aug.—Oct. v. v. Flowers pale yellow. It is an intermediate species between the preceding and *P. alba*.

virgata.

4. P. glabra; caule simplicissimo, foliis omnibus lyrato-sinuatis, racemulis subsecundis, floribus pendulis, calycibus glabris 8-fidis 10-floris.—Mich. fl. amer. 2. p. 83.

P. autumnalis. Walt. fl. car. 193.

In sandy fields, near ditches: New Jersey to Carolina.

24. Aug. Sept. v. v. From three to six feet; flowers pale purple. This plant has a very fine effect in large plantations. Michaux quotes Gronovius; but his *P. autumnalis* belongs to *P. rubicunda*.

simplex.

5. P. caule simplicissimo, foliis superioribus lineari-lanceolatis integerrimis, radicalibus lanceolatis sinuatis, racemo terminali simplicissimo, floribus nutantibus, calycibus sub-8-floris.

P. virgata. Willd. sp. pl. 3. p. 1533.

In Carolina. Enslen. 24. July, Aug. v. s. in Herb. Enslen. About two feet high; flowers purple.