

FRUCTITUBIA SYNANTHERIA. LACTUCEÆ. Hieracium. 63

700 dunense, Vh! 2. 7. In maritimis Flandriæ! Hollandiæ! Ze-
landiæ!

701 umbellatum, L. 2. 7—9. In sylvaticis!

702 lineare = H. umbellatum. β. Pers. 2. 9, 10. In sylvis Verv.
(Lej!)

DREPANIA. Juss. Cass.

703 barbata Desf. (I). 8. In arenosis propè Orroir (Paternost. Hoor!)

ARNOSERIS. Gærtn. Cass.

704 minima = Hyoseris, L. (I). 6, 7. In agris arenosis!

705 foetida = Hyoseris, L. 2. 6. 7. In Flandriâ fide Roucel.
Dub. civ.

Subtrib. 4. SCORZONERÆ. Cass.

HYPOCHÆRIS. Lin

§. 1. PORCELLITES. Cass.

706 maculata, L. 2. 6, 7. In pratis montosis siccis!
β. uniflora.

707 radicata, L. 2. 6—8: In pratis et agg. graminosis!

§. 2. HYPOCHÆROTYPUS. = Hypochæris Gærtn., Scop.

708 glabra, L. (I). 6—8, In arvis arenosis!

HYOSERIS. Juss. Cass.

709 scabra, L. (I). 6, 7. In Hann. Gall. (Hécart!)

LEONTODON = L. et Thrinacia, Dec!

§. 1. THRINCIA. Dec. Cass.

710 hirtum, L. 2. 6—9. In glareosis et ageribus!

711 saxatile, Lam. (I). 7, 8. In arenosis marit!

§. 2. APARGIA. Moench.

712 hispidum, L. 2. 6—9. In pascuis argillosis!

713 squamosum, Lamk. 2. 6. In pascuis montosis Walcourt (Derm!)

714 hastile, L. 2. 5—7. In pascuis montosis. Luxemb. (March!)
β. hirtifolium.

§. 3. SCORZONEROIDES., Moench.

715 autumnale, L. 2. 7—10. In pascuis et versuris!

716 contortum: foliis runcinatis hispidis; scapis uni-bi-floris patulis
squamosis glabris, pericliniis glabris demum contortis. 2. 6—8.
In arenosis marit! — Fol. L. hispidi; scapi L. autumnalis; pe-
riclinii proprii.