

L. DUMICOLA, comb. nov. (*Crepis dumicola* Hiern). ANGOLA: in herb-grown pasturage near Forte P. Amelia, 2063. Florets pale violet.—Angola.

L. ANDONGENSIS Hiern. ANGOLA: in grass-lands along R. Ludira at Cuando, Caconda, after bush-fires, 1728. A perennial herb, stem thick and fistular, branched towards top; lower leaves much larger than those Hiern describes, reaching 23×7 cm.; florets white.—Angola.

SONCHUS BIPONTINI Aschers. ANGOLA: in grass-lands after bush-fires at Camasaio near Malange, 1196; in wet grassy meadows between Forte P. Amelia and R. Cubango, 2187.

Var. PINNATIFIDA Oliv. & Hiern. ANGOLA: here and there in grassy thickets, N'Dallatondo, Cazengo, alt. 750 m., 4375.—(Species) Tropical Africa except West.

S. WELWITSCHII Chiov. (*S. Fischeri* O. Hoffm.). ANGOLA: in sandy exposed situations near the fazenda Palmyra, Cazengo, 479; in open forests after bush-fires, N'Bango near Malange, 1192; in open thickets at Micango, Cuiriri, 4090; skirting road between Cammonda and R. Carunda, Cazengo, 5080.—Angola, Congo, Rhodesia, Uganda.

S. OLERACEUS L. ANGOLA: Vulangombe—N'Golo road, Malange, 1187.—Cosmopolitan weed.

S. ASPER L. ANGOLA: Quizanga, Malange, 1157.—Cosmopolitan weed.

S. ELLIOTIANUS Hiern. ANGOLA: Malange after bush-fires, 1163, 1168; on stony ground in open grassy thickets between Forte P. Amelia and R. Cubango, 1979.—Angola, Congo, Rhodesia, Transvaal.

Sonchus cubangensis S. Moore, sp. nov. *Herba* perennis, glabra, circa sesquispithamea viva ex cl. delectorii glauca in sicco vero brunnescens; *caule* e rhizomate circa 6 mm. diam. orto erecto simplici sat bene folioso; *foliis* (paucis radicalibus confertis) pro genere parvis lanceolatis vel lanceolato-ovatis spinuloso-acuminatis basi amplexicaulibus margine dentatis dentibus induratis junioribus fere integris in bracteas gradatim transeuntibus; *capitulis* circa 16-flosculosis anguste campanulatis in corymbum sublaxum oligocephalum bracteatum foliis longiorem digestis; *pedunculis propriis* capitula subæquantibus excedentibusve; *involuci* phyllis paucis 4-serialibus interioribus sensim longioribus exterioribus ovatis interioribus ovato-oblongis oblongisve omnibus obtusis et microscopice serrulatis nisi integris; *corollis* exsertis dilute flavis; *achæniis* adhuc valde crudis compressis ovato-oblongis truncatis obscure striatis quam pappi setæ a corollis facile superatæ annulatim connatæ multo brevioribus.

Hab. ANGOLA: near Forte P. Amelia, Cubango, 2956.

An erect herb some 30 cm. high. Stem stoutish, prominently ribbed, 4 mm. across below, narrowing to about 3 mm. Radical leaves few, apparently about 2·5 cm. long, but all seen broken off above. Cauline leaves lowest 4·5 cm. long, decreasing to 3-2 cm., 10-7 mm. wide at base, but rapidly narrowing upwards, drying brown like the stem. Inflorescence 12×6 cm., its bracts in shape like the uppermost leaves, gradually diminishing from 15 to 5 mm. in length. Expanded heads 18×12 mm. Outermost involucral leaves 3·5-5 mm. long; intermediate up to 12 mm., innermost to 14 mm. long. Corollas 13 mm., achenes 1·5 mm., pappus 8·5 mm. long.

A very distinct species, recognized by the short amplexicaul foliage and the broad involucral leaves.

CAMPANULACEÆ.

CEPHALOSTIGMA PERROTTETII A. DC. ANGOLA: in the woods and masacallas of Cassuango, Cuiriri (a masacalla is an abandoned *Pennisetum* plantation overgrown with trees), fl. April, 2991.—Western Tropical Africa.

LIGHTFOOTIA COLLOMIOIDES A. DC. ANGOLA: in herb-grown thickets and woods at T'Chirandangombe on the left bank of R. Luassingua, Ganguellas, fl. April, 2702; at Dongo near Cuelai, Massaca, 3460.—Southern Tropical Africa, Uganda.

L. ANNUA A. DC. ANGOLA: in grassy thickets behind my residence at Malange, a late specimen, May, 1435; Yumba, Calulo, Libolo country, 6358.—Angola.

L. MARGINATA A. DC. ANGOLA: in sandy poor ground once cultivated by natives, near Fte. P. Amelia, Cubango, 2389; Cului, 2921.—Southern Tropical and South Africa.

WAHLENBERGIA CALEDONICA Sond. ANGOLA: among *Glumaceæ* in moist situations below the Harta of Fte. P. Amelia, Cubango, fl. Dec., 2342.—Rhodesia, Angola, South Africa.

Cyphia cacondensis R. Good, sp. nov. *Herba* glabra; *stirpe* terete voluta; *foliis* petiolatis ellipticis lanceolatis obtusis basi breviter cuneatis, marginibus calloso-crenatis, utrinque glabris, infra venis conspicuis; *floribus* pedicellatis solitariis in axillis foliorum superiorum, pedicellis supra medium minute bibracteatis; *ovario* semi-supero; *calyce* tubo turbinato striato, lobis lanceolatis basi utrinque dente uno præditis; *corolla* pallide violaceo-cærulea, lobis angustis ligulatis versus apicem crassiusculis intus basi hirsutis; *staminibus* quam lobis corollæ dimidio brevioribus, antheris dorso hirsutissimis; *stigmate* obliquo.

Hab. ANGOLA: in open herb-grown woods between Caconda and Bissava on route, fl. March, 4302.

Rootstock apparently tuberous. Stem climbing in grasses. Leaves up to 6 by 4 cm.; petioles about 5 mm.; pedicels 5 mm.; calyx-