

*Helminthotheca echioides* (L.) Holub, *Folia Geobot. Phytotax.* **8**: 176 (1973).

var. **mollis** (Duby) P. D. Sell comb. nov.

*Helminthia echioides* var. *mollis* Duby, *Bot. Gall.* **1**: 300 (1828).

var. **pratensis** (Chevall.) P. D. Sell comb. nov.

*Helminthia echioides* var. *pratensis* Chevall., *Fl. Gén. Env. Par.* **2**: 352 (1827).

var. **incisa** P. D. Sell var. nov.

Holotype: Garden of 54, Cambridge Road, Impington, Cambs, v.c. 29, 52/436619, 15 July 1998, P. D. Sell no. 98/92 (CGE) (1 plant on 6 sheets).

*Planta* erecta, pilis longioribus rigidis aequae cum brevibus vestita. *Folia* magna, costam versus ad partem dimidiam lobis latis dentatis incisa. *Involucri bractee exteriores* ad basin cordatae.

*Sonchus oleraceus* L., *Sp. Pl.* 794 (1753).

var. **litoralis** P. D. Sell var. nov.

Holotype: Shingle beach between Cley and Salthouse, E. Norfolk, v.c. 27, 63/059451, 21 September 1967, P. D. Sell no. 67/1775 and N. Marchant (CGE).

*Caulis* 10–50 cm. *Folia* saepe rubripurpureotincta. *Capitula* pauca, in fasciculo denso congesta.

*Sonchus asper* (L.) Hill, *Herb. Brit.* **1**: 47 (1769).

var. **sabulosus** P. D. Sell var. nov.

Holotype: Stabilised shingle at back of beach, north of Dunwich, E. Suffolk, v.c. 25, 62/479709, 26 June 1990, P. D. Sell no. 90/157 and J. G. Murrell (CGE).

*Caulis* ad 50 cm. *Folia* plerumque divisa, aculeatissima. *Capitula* pauca, in fasciculo denso congesta.

*Lactuca virosa* L., *Sp. Pl.* 795 (1753).

forma **lactucarii** (Lamotte) P. D. Sell stat. nov.

*L. lactucarii* Lamotte, *Prodr. Fl. Plat. Centr.* **2**: 467 (1881).

*L. scariola* forme *virosa* var. *lactucarii* (Lamotte) Rouy, *Fl. France* **9**: 199 (1905).

*Crepis vesicaria* L., *Sp. Pl.* 805 (1753).

var. **rutilans** (Lacaita) P. D. Sell comb. et stat. nov.

*C. rutilans* Lacaita, *Jour. Linn. Soc. Bot.* **44**: 128 (1918).

*Pilosella* × *stoloniflora* (Waldst. & Kit.) F. W. Sch. & Sch. Bip. in *Flora (Regensb.)* **45**: 423 (1862).

nothosubsp. **schuriana** (Nägeli & Peter) P. D. Sell comb. nov.

*Hieracium* × *stoloniflorum* subsp. *schurianum* Nägeli & Peter, *Hier. Mittel-Europas: Piloselloiden* 326 (1885).

Probabiliter = **P. aurantiaca** subsp. **carpathicola** × **officinarum** subsp. **euronota**

*Hieracium* L.

In Section Sabauda (Fr.) Arv.-Touv.

*Hieracium vagum* Jord., *Cat. Grenoble* 21 (1849).

forma **croceostylum** (Pugsley) P. D. Sell comb. et stat. nov.

*H. croceostylum* Pugsley in *Jour. Linn. Soc. London (Bot.)* **54**: 310 (1948).

***Hieracium prominentidens*** P. D. Sell nom. et stat. nov.

*H. dispalatum* var. *macrodon* Sudre, *Hier. Centr. France* 24 (1902), non *H. macrodon* Nägeli & Peter, *Hier. Mittel-Europas: Archieracien* 84 (1886).

*Hieracium sabaudum* L., *Sp. Pl.* 804 (1753).

forma **bladonii** (Pugsley) P. D. Sell comb. et stat. nov.

*H. bladonii* Pugsley in *Jour. Linn. Soc. London (Bot.)* **54**: 299 (1948).

In Section Hieracioides Dumort.

*Hieracium umbellatum* L., *Sp. Pl.* 804 (1753).

var. **sarniense** P. D. Sell var. nov.

Holotype: Granite cliffs, north coast of Jersey, Channel Islands, 17 July 1901, L. V. Lester-Garland as E. F. & W. R. Linton, *Set of British Hieracia* no. 185 (CGE).

*Caulis* 10–30 cm, robustus, succulentus. *Folia* crassa, succulenta, anguste vel late elliptica, ad apicem obtusa, inferiora ad c. 5 cm lata, prope caulis basin fasciculata, sursum minora.

var. **latifolium** (W. R. Linton) P. D. Sell stat. nov.

*H. umbellatum* forma *latifolium* W. R. Linton, *Brit. Hier.* 98 (1905).

In Section Foliosa (Fr.) Arv.-Touv.

***Hieracium reticulatiforme*** P. D. Sell nom. et stat. nov.

*H. reticulatum* var. *angustifolium* Pugsley in *Jour. Linn. Soc. London (Bot.)* **54**: 280 (1948), non *H. angustifolium* Hoppe, *Bot. Taschenb.* **1799**: 129 (1799).

***Hieracium reayense*** (Pugsley) P. D. Sell stat. nov.

*H. reticulatum* var. *reayense* Pugsley in *Jour. Linn. Soc. London (Bot.)* **54**: 280 (1948).

***Hieracium melanoglochis*** (E. F. Linton) P. D. Sell stat. nov.

*H. corymbosum* var. *melanoglochis* E. F. Linton in W. R. Linton, *Brit. Hier.* 88 (1905).

In Section Tridentata (Fr.) Arv.-Touv.

***Hieracium charitodon*** P. D. Sell sp. nov.

Holotype: Waterfall at the head of Glyn Tarell, Brecon, v.c. 42, 25 July 1900, A. Ley (CGE).

*Herba* perennis aphyllopoda vel hypophyllopora, caudice ramoso. *Caulis* 35–100 cm, pallide luteoviridis, ad basin saepe purpurascens, gracilis vel robustus, infra pilis simplicibus eglandulosis densis longis pallidis, sursum paucioribus brevioribusque, et in parte superiore pilis stellatis numerosis vestitus. *Folia* in pagina superiore pallide vel mediocriter viridia, in pagina inferiore pallidiora, 5–15, sursum gradatim decrescentia; folia basalia et caulina infera lamina 6–11 × 1.5–4.0 cm, anguste elliptica vel oblongoelliptica vel lanceolatooblonga, ad apicem obtusomucronatum vel acutum gradatim angustata, dentibus mammiformibus vel longiusculis angustisque dentata, ad basin in petiolum alatum ad 4 cm longum attenuata; caulina mediana lamina 4–9 × 1.5–4.0 cm, linearielliptica