

*Sonchus canariae* Pit. in Pit. & Pr., Iles Canaries, Fl.Arch.p.261; 1908  
f. *angustissimus* Kunkel, forma nova

A forma typica foliis linear-lanceolatis, integris vel leviter denticulatis differt.

HOLOTYPUS: Kunkel 11018 (G.), 8.V.1967; Gran Canaria, Cenobio de Valerón 180 m.; very rare.

*Sonchus bourgeaui* Sch. Bip.

Known from near Mogadar (Maroc), La Graciosa, Lanzarote and northern Fuerteventura; herewith a new locality of this species: Fuerteventura, -- Jandía, Pico de la Zarza, 780 m. (Ku. 12545).

*Sonchus pinnatifidus* Cav., Anal. Cienc. Nat. 4: 78; 1801

var. *integrifolius* Kunkel, var. nov.

Frutex 1,5 - 2,5 m. alt.; a var. *pinnatifidus* folia integerrima differt. Lamina ovaliforme-elliptica, usque ad 20 cm. longa et 10 cm. lata; margine leviter undulatae.

HOLOTYPUS: Kunkel 18831 (G.), 22.II.1976; Lanzarote, Malpaís de la Corona, 140 m.; frequent, on rocks.

*Taeckholmia capillaris* (Svent.) Boulos

Material (Ku. 15595) from Arteana. 300 m. (near Fataga, Gran Canaria) should belong to this taxon; however a revision with *T. microcarpa* Boulos, from Tenerife, is recommended.

*Taeckholmia regis-jubae* (Pit.) Boulos

In case this rather variable taxon remains separated from related species, a new locality may be added: Tenerife, El Sauzal, 350 m.; Ku. 14582. The species was known before to be endemic in La Palma, Gomera, and Gran Canaria.