

- pinnatus.* 8. *S. pedunculis nudis, calycibus lævibus, foliis pinnatis; pinnis lineari-lanceolatis subdentatis.*  
Wing-leav'd Sow-thistle.  
*Nat. of Madeira. Mr. Francis Masson.*  
*Introd. 1777.*  
*Fl.* G. H. 5.
- radicatus.* 9. *S. pedunculis nudis calycibusque glabris, caule subnudo, foliis radicalibus lyratis utrinque lævibus lobis triangulari-ovatis.*  
Long-rooted Sow-thistle.  
*Nat. of the Canary Islands. Mr. Francis Masson.*  
*Introd. 1780.*  
*Fl. July.* G. H. 5.
- floridanus.* 10. *S. pedunculis squamosis, foliis lyrato-hastatis.* *Syst. veget. 712.*  
Small-flower'd Sow-thistle.  
*Nat. of North America.*  
*Cult. 1713. Philosoph. trans. n. 337. p. 182. n. 20.*  
*Fl. July.* H. 4.
- sibiricus.* 11. *S. pedunculis squamatis, foliis lanceolatis indivisis sessilibus.* *Sp. pl. 1118.*  
Willow-leav'd Sow-thistle.  
*Nat. of Sweden and Russia.*  
*Cult. 1759, by Mr. Philip Miller.*  
*Fl. July and August.* H. 4.
- tataricus.* 12. *S. pedunculis nudis, foliis lanceolatis dentatis runcinatis.* *Syst. veget. 712.*  
Tartarian Sow-thistle.  
*Nat. of Siberia.*  
*Introd. 1784, by William Pitcairn, M. D.*  
*Fl.* H. 4.  
13. S.