

On road sides, in shady woods: Virginia and Carolina.
♂. July—Sept. v. v. Flowers very small, blue.
This plant has been used for curing the bite of the
rattlesnake, in the same manner as *Prenanthes Ser-*
pentaria, and is known by the name of *Gall of the*
Earth.

- acuminatus.* 7. *S. pedunculis subsquamosis, floribus paniculatis, foliis*
radicalibus subruncinatis, caulinis ovatis acuminatis
petiolatis medio denticulatis. —*Willd. sp. pl. 3.*
p. 1521.

Lactuca villosa. Jacq. hort. schœnb. 3. t. 367?

In low shady places: Pennsylvania to Virginia. ♂. Aug.
Sept. v. v. Flowers small, blue.

- pallidus.* 8. *S. racemo composito terminali, foliis lanceolato-ensi-*
formibus amplexicaulibus dentatis. *Willd. sp. pl. 3.*
p. 1521.

Lactuca canadensis. Sp. pl. 1119.

Icon. Rob. ic. 148 & 151.

On road sides and in woods: Canada to New England.
♀. July—Sept. v. v. Flowers small, yellow.

- pulchellus.* 9. *S. pedunculis squamosis, floribus corymboso-racemosis,*
foliis caulinis cordato-amplexicaulibus ovato-oblongis
acutis integerrimis glabris.

On the banks of the Missouri. ♀. Sept. v. s. Flowers
large, beautiful blue.

601. HIERACIUM. Gen. pl. 1238.

- pusillum.* 1. *H. pusillum, villosissimum; caule simplici erecto uni-*
floro remote subbisfoliato, foliis lanceolatis attenuatis
acutis integerrimis, caulinis linearibus, calyce villosis-
simo. —*Willd. sp. pl. 3. 1561.*

In Labrador. *Colmaster.* ♀. July. v. s. in *Herb.*
Dickson. A very small plant; flowers, as in all the
following species, yellow, the size of a common daisy.
It strongly resembles *H. alpinum*, and is probably only
a diminutive variety thereof.

- venosum.* 2. *H. scapo nudo corymboso-paniculato glabro, pedicellis*
filiiformibus, foliis obovato-lanceolatis supra rarer pil-
losis, subtus nudis margine ciliatis glanduloso-denticu-
latis, venis coloratis, calycibus glabris. —*Willd. sp.*
pl. 3. p. 1570.

In shady fertile woods: Canada to Carolina. ♀. June
—Aug. v. v. Flowers middle size, of a beautiful