

- pinnatus.* 8. *S. pedunculis nudis, calycibus lævibus, foliis pinnatis, pinnis lineari-lanceolatis subdentatis.*
Wing-leav'd Sow-thistle.
Nat. of Madeira. Mr. Francis Masson.
Introd. 1777.
Fl. G. H. 5.
- radicatus.* 9. *S. pedunculis nudis calycibusque glabris, caule subnudo, foliis radicalibus lyratis utrinque lævibus lobis triangulari-ovatis.*
Long-rooted Sow-thistle.
Nat. of the Canary Islands. Mr. Francis Masson.
Introd. 1780.
Fl. July. G. H. 5.
- floridanus.* 10. *S. pedunculis squamosis, foliis lyrato-hastatis.* *Syst. veget. 712.*
Small-flower'd Sow-thistle.
Nat. of North America.
Cult. 1713. Philosoph. trans. n. 337. p. 182. n. 20.
Fl. July. H. 4.
- sibiricus.* 11. *S. pedunculis squamatis, foliis lanceolatis indivisis sessilibus.* *Sp. pl. 1118.*
Willow-leav'd Sow-thistle.
Nat. of Sweden and Russia.
Cult. 1759, by Mr. Philip Miller.
Fl. July and August. H. 4.
- tataricus.* 12. *S. pedunculis nudis, foliis lanceolatis dentatis runcinatis.* *Syst. veget. 712.*
Tartarian Sow-thistle.
Nat. of Siberia.
Introd. 1784, by William Pitcairn, M.D.
Fl. H. 4.