

**Xizang (西藏):** Yadong (亚东), in fruticeto in declivio, alt. 3100 m, 1974.09.15., Qinghai-Xizang Exped. 74-2566 (Typus, PE), 74-2216; Nyalam (聂拉木), ad marginem sylvae, alt. 3300—3400m, Xizang Exped. Pl. Med. (西藏中草药队) 1533; Bomi (波密), in sylva ad declivium, alt. 2400 m, Z. C. Ni (倪志诚) 1604.

## 2. 栉齿细莴苣 新种

***Stenosaris triflora*** Chang et Shih, sp. nov.

Affinis *Ch. graciliflorae* (Wall. ex DC.) Shih, a qua pectinibus auriculiformibus ad rachides foliorum bene differt.

Herba perennis. Caulis erectus, 0.5—2.5 m altus, superne paniculato-ramosus, ramis gracilibus, pilis multicellularibus noduliformibus sparse vel dense obiectus. Folia inferiora ignota; folia media lyrato-pinnatisecta, lobis terminalibus ovato-orbiculiformibus, -cordatis vel lyrato-lanceolatis, 9—11 cm longis, 7—9 cm latis, apice acuminatis, lobis latelaribus 2-jugis, ovatis vel ellipticis, 5.5—7 cm longis, 2.5—4.5 cm latis, apice acuminatis vel acutis, omnibus margine mucronulato-dentatis vel-denticulatis, rhachidibus auriculiforme pectiniformibus; folia superiora mediis similia, sed minora; omnia textura tenuia, utrinque strigulosa. Cacathidia numerosa, adapices caulium et ramorum in inflorescentiam paniculatam terminalem disposita. Involucrum anguste cylindricum, 1 cm longum, 1—1.5 mm diam., calyculatum, phyllis 1-seriatis, lanceolatis vel lineari-lanceolatis, 1 cm longis, 2 mm latis, apice obtusis vel acutis. Flosculi ligulati 3, purpurei. Achaenia longe elliptica, 3.5 mm longa, 1mm lata, leviter compressa, facie costulis 4—5, margine incrassata, utrinque pilis brevissimis vel valde obscuris in series transversas unduatas dispositis obiecta, apice in rostrum breviter contracta, rostris 1 mm longis. Pappi setae extimae e pilis brevissimis, obscuris, sparsis constantes; pappi setae interiores e pilis numerosis, tenuibus, 6 mm longis, fragilibus efformantes; omnes pallide albo-flavescentes.

**Yunnan (云南):** Fugong (福贡), alt. 2600m, 1933.10.10., H. T. Tsai (蔡希陶) 54364 (Typus, PE), 58675, 59100; Jinping (金屏), in locis humectis in sylva, alt. 1700m, 1951.08.15., P. Y. Mao (毛品一) 358; Luchun (绿春), ad marginem sylvae in declivio umbroso, alt. 2100 m, 1973.10.17., D. D. Tao (陶德定) 887.

## 3. 细莴苣 图 1:7

***Stenosaris graciliflora*** (Wall. ex DC.) Shih, comb. nov. — *Lactuca graciliflora* Wall. ex DC., Prodr. 7: 139, 1838; Hook. f., Fl. Brit. Ind. 3: 406, 1881; S. Y. Hu in Quart. Journ. Taiwan Mus. 20 (1—2): 20, 1967, p. p. — *Prenanthes brunnoniana* auct. non Wall. ex DC.: 西藏植物志 4: 957, 1985, p.p.

四川: 方文培 15027; 青藏队 83-14344; H. Smith 10972. 云南: 王启无 69082; 冯国楣 3290, 7825, 2117, 3290; 秦仁昌 24009, 31077, 39627; 俞德浚 20765, 23156; 昆明所中甸队 1982; 赵致光 30627. 西藏: 青藏队补点组 751291; 西藏中草药普查队 1533; 青藏队 82-10443, 741980, 74-2837; 青藏队植被组 5312; 青藏队草原组 6965.